

A California Toolkit to Transform Maternity Care

Obstetric Hemorrhage Toolkit

HOSPITAL LEVEL IMPLEMENTATION GUIDE

THIS COLLABORATIVE PROJECT WAS DEVELOPED BY:

THE OBSTETRIC HEMORRHAGE TASK FORCE

THE MATERNAL QUALITY IMPROVEMENT PANEL

CALIFORNIA MATERNAL QUALITY CARE COLLABORATIVE

MATERNAL, CHILD AND ADOLESCENT HEALTH DIVISION; CENTER FOR FAMILY HEALTH

CALIFORNIA DEPARTMENT OF PUBLIC HEALTH



CMQCC OBSTETRIC HEMORRHAGE TOOLKIT HOSPITAL LEVEL IMPLEMENTATION GUIDE

Debra Bingham, DrPH, RN^a, Kathryn Melsop, MS^a, Elliott Main, MD^{a,b}

California Maternal Quality Care Collaborative (CMQCC)^a and The California Pacific Medical Center (CPMC)^b

Suggested citation:

Bingham, D., Melsop, K., Main, E. (2010). CMQCC Obstetric Hemorrhage Hospital Level Implementation Guide. The California Maternal Quality Care Collaborative (CMQCC). Stanford University, Palo Alto, CA.

Funding for the development of this toolkit was provided by:

Federal Title V Funding and the California Department of Public Health; Maternal, Child and Adolescent Health Division, The California Perinatal Quality Care Collaborative, and Stanford University.

CMQCC OB Hemorrhage Toolkit Hospital Level Implementation Guide is part of the Transforming Maternity Care Series and was developed under contract #08-85012 between the California Department of Public Health: Maternal, Child and Adolescent Health Division and the California Maternal Quality Care Collaborative.

Copyright: This document is in the public domain and may be used and reprinted without permission except those copyrighted materials noted for which further reproduction is prohibited without the specific permission of copyright holders.

For correspondence, please contact

CMQCC

Kathryn Melsop, MS
Managing Editor, Transforming Maternity Care Series
Medical School Office Building, Stanford University
251 Campus Drive
Stanford, CA 93405
Phone: (650) 725-6108
FAX: (650) 721-5751
email: melsop@cmqcc.org
Website: <http://www.cmqcc.org/>

CA Department of Public Health

Connie Mitchell, MD, MPH
1615 Capitol Avenue
PO Box 997420, MS 8306
Sacramento, CA 95899-7420
Phone: (916) 650-0327
email: connie.mitchell@cdph.ca.gov
Website: <http://www.cdph.ca.gov>

ACKNOWLEDGEMENTS

CMQCC would like to thank the California Department of Public Health; Maternal Child Health Division leaders Shabbir Ahmad, PhD and Connie Mitchell, MD, MPH.

CMQCC would like to thank volunteer members of the Hemorrhage Task Force (HTF) and the Task Force Co-Chairs, David Lagrew, MD, and Audrey Lyndon, PhD, RND, CNS and the Maternal Quality Improvement Panel (MQIP) for their contributions to the Obstetric Hemorrhage Care Guidelines, Compendium of Best Practices and Hospital Level Implementation Guide.

HEMORRHAGE TASK FORCE

Co-Chairs:

- Audrey Lyndon PhD, RNC, CNS – University of California, San Francisco
- David Lagrew, MD – Saddleback Memorial Hospital

- * Leslie Casper, MD – Kaiser, Southern CA, San Diego Medical Center
- * Nancy Corbett, BSN, RN – Kaiser, Northern CA
- * Maurice Druzin, MD – Stanford University
- * Sue Faron, MN, RNC, CNS – Sharp Mary Birch Hospital for Women
- * Kim Gregory, MD, MPH – Cedars-Sinai Medical Center
- * Andrew Hull, MD, FRCOG, FACOG – University of California, San Diego
- * Valerie Huwe, BSN, RNC – Stanford Hospital, El Camino Hospital
- * Richard Lee, MD – USC, Women’s and Children’s Hospital
- * Holli Mason, MD – Harbor UCLA Medical Center
- * Elliott Main, MD – California Pacific Medical Center
- * Jennifer McNulty, MD – Long Beach Memorial Medical Center, UC Irvine
- * Suellen Miller, PhD, MHA, CNM - University of California, San Francisco
- * Connie Mitchell, MD – MCAH, CA Department of Public Health
- * Mark Rosen, MD – University of California, San Francisco
- * Diana Ramos, MD, MPH – MCAH, County of Los Angeles Public Health
- * Larry Shields, MD – Central Coast Maternal Fetal Medicine
- * Jean-Claude Veille, MD – Sutter Health, Sacramento

MATERNAL QUALITY IMPROVEMENT PANEL

- * Karyn Almryde, MSN, RN – Sharp Mary Birch Hospital for Women
- * James Byrne, MD – Santa Clara Valley Medical Center
- * Mary Campbell Bliss, MS, RN, CNS – Sutter Memorial Sacramento-Sierra Region
- * Brenda Chagolla, MSN, RNC, CNS – Catholic Healthcare West
- * Holly Kennedy, PhD, CNM, FACNM – University of California, San Francisco
- * David Lagrew, MD – Saddleback Memorial Medical Center
- * Audrey Lyndon, PhD, RNC, CNS – University of California, San Francisco
- * Connie Mitchell, MD, MPH – MCAH, CA Department of Public Health
- * Barbara Murphy, MSN, RNC – CPQCC
- * Karen Ramstrom, DO, MSPH – MCAH, CA Department of Public Health
- * Linda Walsh, PhD, CNM, FACNM – University of San Francisco

OB HEMORRHAGE TOOLKIT

EXECUTIVE COMMITTEE

- Berneva Adams, MD – Southern California Kaiser Permanente Medical Group
- Shabbir Ahmad, DVM, MS, PhD – MCAH, California Department of Public Health
- Lisa Bollman, RNC, MSN, CPHQ – Community Perinatal Network
- James Byrne, MD – Santa Clara Valley Medical Center
- Leslie Cragin, CNM, PhD, PhD, FACNM – San Francisco General Hospital Medical Center
- Arlene Cullum, MPH – Sutter Medical Center, Sacramento
- Suzanne Delbanco, PhD – Arrowsight, Inc.
- Tracy Flanagan, MD – Kaiser Permanent Medical Group, Richmond
- Bill Gilbert, MD – Sutter Health, Sacramento Sierra Region
- Jeffrey Gould, MD, MPH – CPQCC, Stanford University
- Lisa Korst, MD, PhD – Childbirth Associates, LLC
- Leslie Kowalewski – March of Dimes
- John Kriege – Office of Statewide Health Planning and Development
- David Lagrew, MD – Saddleback Memorial Medical Center
- Elizabeth Lawton, MHS – MCAH, California Department of Public Health
- Michael Lu, MD, MPH – University of California, Los Angeles
- Audrey Lyndon PhD, RNC, CNS – University of California, San Francisco
- Kathryn Melsop, MS – CMQCC
- Connie Mitchell, MD, MPH – MCAH, CA Department of Public Health
- Christine Morton, PhD – CMQCC
- Barbara Murphy, RNC, MSN – CPQCC, Stanford University
- Alan Oppenheim, MA, MPH – CDHS, Center for Health Statistics
- Steven Parry, MD, FACOG, Medical Benefits Branch, MCBWARD
- Manuel Porto, MD – University of California, Irvine
- Diana Ramos, MD, MPH – County of Los Angeles Public Health
- Karen Ramstrom, DO, MSPH – MCAH, California Department of Public Health
- Usha Ranji, MS – Kaiser Family Foundation
- Leona Shields, PHN, MN, CNP – MCAH, California Department of Public Health
- Randy Winter, MD – Doctors Medical Center of Modesto
- Teresa Wray, MD – Kaiser Permanente South

CMQCC OBSTETRIC HEMORRHAGE QUALITY IMPROVEMENT COLLABORATIVE EXPERT PANEL

- David Lagrew, MD – Saddleback Memorial Medical Center
- Bev VanderWal, CNS – Long Beach Memorial Medical Center
- Julie Arafah, RN, MSN – Stanford University
- Holli Mason, MD – Harbor UCLA Medical Center
- Mark Rollins, MD – University of San Francisco, California

CMQCC

- * Elliott Main, MD, Principal Investigator
- * Debra Bingham, RN, DrPH, Executive Director
- * Kathryn Melsop, MS, Program Manager
- * Christine Morton, PhD, Program Manager
- * Valerie Cape, Program Administrator

OBSTETRIC HEMORRHAGE TOOLKIT HOSPITAL LEVEL IMPLEMENTATION GUIDE TABLE OF CONTENTS

Introduction	98
Composite Case	99
Obstetric Hemorrhage Task Force	100
Getting Started: Model for Improvement.....	101
Rapid Cycle QI Method: MAP-IT	
Mobilize	102
Assess.....	103
Plan	107
Implement	108
Track	112
Appendices	
Appendix A: Sample Obstetric Hemorrhage Policy and Procedure.....	114
Appendix B: Obstetric Hemorrhage Care Guidelines: Checklist.....	121
Appendix C: Obstetric Hemorrhage Care Guidelines: Flowchart	125
Appendix D: Obstetric Hemorrhage Care Guidelines: Table Chart	126
Appendix E: Obstetric Hemorrhage Quality Improvement Collaborative.....	127
Appendix E.1: Risk Assessment Audit Tool	128
Appendix E.2: Active Management of Third Stage Labor Audit Tool.....	129
Appendix E.3: Methods for Developing Training and Tools for Quantitative Blood Loss Measurement	130
Appendix E.4: Cumulative Blood Loss Measurement Audit Tool	131
Appendix E.5: Team Debrief Form	132
Appendix F: MAP-IT Worksheet, Sample Worksheet.....	133
Appendix G: Obstetric Hemorrhage Quality Improvement Measurement Grid.....	135

INTRODUCTION

CALIFORNIA MATERNAL QUALITY CARE COLLABORATIVE (CMQCC)

CMQCC's mission is to transform maternity care in California to end preventable death and injury. To achieve this we will:

- Define and implement best practices for public health, communities and women with quality, safety and social justice as the clear priorities of every decision and action.
- Promote communication and collaboration between all maternity stakeholders.
- Gather, review and organize maternity data and statistics into actionable information.
- Build the next generation of maternal health leaders to continue the growth and scope of CMQCC.

We are devoted to eliminating preventable maternal death and injury and promoting equitable maternity care in California by bringing resources, tools, measures and quality improvement techniques to providers, administrators and public health leaders. This is a long-term collaborative effort of many organizations and individuals with Title V funding from the California Department of Public Health (Maternal, Child and Adolescent Health Program) and the California Perinatal Quality Care Collaborative (CPQCC). We are over 250 clinicians, public health leaders, key payers and representatives of the public all devoted to improving childbirth outcomes.

For more information about CMQCC, visit our website at www.cmqcc.org.

The Obstetric Hemorrhage Toolkit, Hospital-level Implementation Guide was reviewed by the California Department of Public Health, Maternal, Child and Adolescent Health Division and is a resource, but doesn't define the standard of care in California. Readers are advised to adapt the guidelines and toolkit based on their local facility's level of care and patient populations and are not to rely solely on guidelines presented here.

IMPACT OF OBSTETRIC HEMORRHAGE

The rate of maternal deaths has nearly tripled from 6 per 100,000 in 1996 to 17 per 100,000 annual births in 2006.⁽¹⁾ Alarming, the rate for African American women has risen from 27.7 to 45.7 per 100,000 live births between 1999 and 2006.⁽¹⁾ Obstetric hemorrhage is one of the leading causes for maternal death and is a major cause of maternal morbidity. In 1997, 2.4% of all live births in California were complicated by postpartum hemorrhage.⁽²⁾ Nationwide, blood transfusions increased 92% during delivery hospitalizations between 1997 and 2005.⁽³⁾

COMPOSITE CASE

The following is an outline of a composite case (combined elements from multiple cases with indentifying features, including all person identifiers, removed to ensure patient confidentiality) that demonstrates how a normal low-risk pregnancy and birth can at times quickly escalate to an emergent situation and death. An outline of learning points and opportunities for quality improvement (QI) opportunities based on the case follows.

Composite Case Example: A 24yo woman, G2 P1 at 38 wks gestation was induced for “tired of being pregnant”:

- After 8hr active phase and 2hr 2nd stage, she gave birth (NSVD) to an 8lb 6oz infant.
- After placental delivery, she had an episode of atony that firmed with massage. A second episode of atony responded to IM methergine and the physician went home (now 1am).
- The nurses called the physician 30min later to report more bleeding and further methergine was ordered.
- 60min after the call, the physician performed a D&C with minimal return of tissue. The woman received more methergine.
- 45min later a second D&C was performed, again with minimal returns. EBL at this point >2,000 ml.
- Delays in blood transfusion occurred because of inability to find proper tubing.
- Anesthesia was delayed, but a second IV started for more crystalloid. VS became markedly abnormal, P=144, BP 80/30.
- One further methergine given and patient taken for a 3rd D&C. She received 2u PRBCs by this point.
- After D&C is complete, she had a cardiac arrest from hypovolemia/hypoxia and was taken to the ICU where she succumbed 3 hours later.

QI Opportunities and Learning Points from the above composite case: How to reducing Mortality and Morbidity from OB Hemorrhage?

- Need a medical indication before performing an induction
- No documentation of actual blood loss, e.g., *what does “more bleeding” mean?*
- Only a few treatments tried, e.g., Methergine and D&C, and then repeated, even when they were ineffective
- Underestimation of blood loss
- Delay in administration of blood
- Lack of working equipment
- Delay in response from other team members
- Delays in adequate resuscitation

- Lack of an organized standardized team approach

OBSTETRIC HEMORRHAGE TASK FORCE

DEVELOPMENT OF BEST PRACTICES, TOOLS, CARE GUIDELINES

The Obstetric Hemorrhage Task Force Tools and Best Practices were developed by a volunteer group of California-based leaders and experts in obstetrics including obstetricians, nurses and midwives under the direction of the California Maternal Quality Care Collaborative's (CMQCC) Maternal Quality Improvement Panel (MQIP). This implementation guide was developed to support local leaders' efforts to successfully implement the best practices and tools for obstetric hemorrhage into practice and to create active quality improvement processes to drive implementation.

The implementation guide is organized to address four broad objectives as described by the Hemorrhage Task Force:

1. Improve ***readiness*** to respond to an obstetric hemorrhage by implementing standardized policies and procedures (general and massive) and developing obstetric hemorrhage rapid response teams.
2. Improve ***recognition*** of OB hemorrhage by performing on-going objective quantification of actual blood loss and triggers of maternal deterioration during and after all births.
3. Improve ***response*** to hemorrhage by performing regular on-site inter-professional hemorrhage drills.
4. Improve ***reporting*** of OB hemorrhage by standardizing definitions and consistency in coding and reporting.

The following elements are critical to affecting change to ensure that improvements are adopted and sustained over time:

- **Leadership**
 - Identifying Leader and Clinician behaviors, including "Champions"
 - Defining the problem and making the case for change
 - Setting goals
 - Allocating resources
- **Policy & Procedure**
 - Agreeing on a plan
 - Creating consistency between departments to improve teamwork and cohesive quality of care; see Appendix for sample policy and protocol
- **Monitoring**
 - Creating audit tools that work; see Appendix for sample tools
 - Communicating progress toward goals

GETTING STARTED

MODEL FOR IMPROVEMENT

The goal of this implementation guide is to provide a simple step-by-step guide for creating quality improvement (QI) changes in your facility to improve care processes associated with obstetric hemorrhage. MAP-IT (Mobilize, Assess, Plan, Implement, Track)—a rapid cycle QI method for outlining change—is presented.(4)

RAPID CYCLE QI METHODOLOGY

MOBILIZE, ASSESS, PLAN, IMPLEMENT, TRACK (MAP-IT)

- **Step 1**
Mobilize Quality Improvement Team
Recruit champions: clinical staff who visualize the ideal, set goals and follow through to realize defined aims

- **Step 2**
Assess the Situation
Determine current practices for response to hemorrhage; identify **QI Data—process and outcome measures**— that inform baseline assessment, guide appropriate implementation, and provide data sufficient to track progress (see Fishbone Diagram, Pareto Chart, Process Flow Maps below)

- **Step 3**
Plan Change Strategies and Tactics
Policy, Scheduling Process, Empowered Physician Leadership: Implement relevant policies and procedures (P&P) or changes to P&P (e.g., ongoing quantitative measurement of blood loss, drills and debriefs) that support a protocol to respond quickly and effectively to obstetric hemorrhage

- **Step 4**
Implement Strategies and Tactics
Conduct **Clinician Education** and training about ongoing quantitative measurement of blood loss, drills and debriefs; education drives convergent department culture, supports buy-in and contributes to successful rollout of change tactics to accomplish the goal

- **Step 5**
Track Progress
Analyze data (see Audit Tools in Appendix) and present results to clinical staff via **Trend Charts** about obstetric hemorrhage (e.g, transfusions, massive transfusions). Review and repeat steps; when necessary, revise newly implemented tactics to ensure sustainable results

Figure 1: MAP-IT QI Methodology



GETTING STARTED

Mobilize

STEP 1. MOBILIZE QUALITY IMPROVEMENT TEAM

Essential team members include individuals with the following roles or skills:

- System Leadership
- Technical Expertise
- Day-to-Day Leadership

In obstetric service units, doctor and nurse leaders would fulfill these essential roles and skills. Utilizing the expertise and tacit knowledge of the front-line leaders and personnel, including nurses, general obstetricians, unit clerks, blood bank technicians is critical to successful implementation. A first step toward designating a team is identifying a system or administrative leader who leads MAP-IT cycles; e.g., meetings held, data collected, assignments completed, monitoring occurred. The following is a sample draft form to document your team:

The Obstetric (OB) Hemorrhage System or Administrative Leader for our birthing facility is:

The nurse leader is: _____

The physician leader is: _____

The members of the team that will work with the system/administrative, nurse, and physician leaders to improve readiness, recognition, response, and reporting are:

NAME	TITLE	eMAIL ADDRESS

Assess**STEP 2. ASSESS THE SITUATION**

In order to identify and prioritize quality improvement needs and next steps, teams can use these and other tools and methods:

- **Fishbone Cause and Effect Diagrams:** A team analysis of how individual process “problem areas” interact as a “whole”(5-7)
- **Process Flowcharts:** A team analysis of the numerous steps needed to complete a process. For example, the number of tasks a nurse must perform in order to obtain emergency medications(5, 7)
- **Failure Modes and Effects Analysis (FMEA):** A systematic analysis of the potential ways in which a failure, e.g., delay in recognition of excessive treatment and delay in adequate treatment of a woman bleeding will occur. The FMEA includes classification by severity or the “effects” of the failure based on the “mode” or type of failure that could occur.(5)
- **Pareto Analysis and Diagrams:** A method to quantify and rank which of the identified opportunities for improvement are the most problematic. The Pareto Principle is the 80/20 rule, e.g., 80% of the failures are caused by only 20% of the processes.(5)

Examples of the Fishbone and Pareto tools are described below.

A. Fishbone Cause and Effect Diagrams:

Quality improvement begins by looking at the “Big Picture” then narrowing down the problem to specific tasks that are reasonable and achievable starting points for change. The fishbone diagram is a common tool for seeing both elements—the whole and its parts.(5) The head of the fish points to the project goal and six (or more) bones extend from the side to capture the inter-relationships between PEOPLE, PROCEDURES, EQUIPMENT, MATERIALS, ENVIRONMENT AND MISCELLANEOUS factors that contribute to the problem. Fishbone diagrams are a good way to get group feedback and insights into the root cause of a problem.

Example:

Use a Fishbone Diagram (Figures 2a, 2b) to identify areas for Quality Improvement using OB Hemorrhage Care Guidelines Checklist, starting with “Pre-Admission” and working through Stages of Hemorrhage to identify problems.

Diagrams can be as extensive and detailed as needed; alternately, diagrams can be drafted for each stage of hemorrhage; e.g., create one diagram that identifies processes of care surrounding Stage 1 Hemorrhage; create a second diagram that identifies processes of care surrounding Stage 2 Hemorrhage, etc.

First drafts can be simple, hand-written versions that are edited, honed and vetted within leadership teams and front line staff to finalize plans for changes.

Labor and Delivery leaders and staff identify processes of care in need of improvement by filling in the “fishbone diagram” based on a step-by-step review of the OB Hemorrhage Care Guidelines Checklist (Figures 2a, 2b). This approach is a useful framework to:

- Identify and define clinical roles and assignments of responsibility during each stage (People, Management)
- Identify medications—usage and dosage—during each stage of hemorrhage; assess that all Obstetric Services clinicians at all levels know and understand medications, usage and dosage (Materials, Process); assess availability and ease of access to critical medication supplies.
- Identify procedures, such as fundal massage, intrauterine balloon, B-Lynch suturing, quantification of blood loss during each stage of hemorrhage (Process, Materials, Equipment)
- Identify blood products and blood replacement during each stage of hemorrhage (Process, Equipment, Materials, Environment); identify availability and ease of access to these products and the supplies needed to administer them.
- Identify procedures for communication between all areas of obstetric care including, but not limited to, communications with the Blood Bank.

Figure 2a.





Figure 2b.



B. Pareto Analysis and Diagrams

Pareto analysis, developed by Italian economist Vilfredo Pareto, is a statistical technique in decision-making to select limited tasks that will have a significant overall effect. The technique is based on the “Pareto Principle” or the “80/20” rule, which states, in quality improvement terms, that most (80%) of the problems are caused by a few (20%) key causes. There are many applications of the Pareto Principal in quality control including the Pareto Diagram, a key tool in Six Sigma (Motorola’s popular business management strategy).(5)

Example: *Use Pareto Analysis to create a Pareto Diagram (Figure 3) to identify areas for Quality Improvement using OB Hemorrhage Care Guidelines Checklist.*

Use the following steps (8) for completing a Pareto Analysis:


- Form a table (e.g., in a spreadsheet program) listing key causes of quality problems and their frequency as a percentage.
- Arrange the rows in the decreasing order of importance of the causes, i.e. the most important problems first.
- Add a cumulative percentage column to the table.
- Plot with causes on x-axis and cumulative percentage on y-axis.
- Join the above points to form a curve.
- Plot (on the same graph) a bar graph with causes on x-axis and percent frequency on y-axis.

- Draw a line at 80% on y-axis parallel to x-axis. Then drop the line at the point of intersection with the curve on x-axis (not shown). This point on the x-axis separates higher priority problems on the left of the line; to the right of the line are problems to be addressed after the higher priority items.

Example:

Use Pareto Analysis to create a Pareto Diagram (Figure 3) to Identify areas for Quality Improvement using OB Hemorrhage Care Guidelines Checklist. Note that in this example, a line drawn on the y-axis at 80% and then dropped down to intersect with the curve would bisect the “Carts” bar. Lack of ongoing quantification of blood loss and delayed availability of blood products become the higher priority issues to be addressed, followed by equipping carts for hemorrhage and fostering education and empowerment for staff about activating a massive hemorrhage protocol.

Figure 3.



Plan

STEP 3. DEVELOP A PLAN TO MEET YOUR PROJECT AIMS

An essential component of a plan is to develop aims or project objectives. The aims or project objectives include the use of specific terms, numerical goals and a time frame or deadline. Teams should work toward consensus agreement for the aims for your facility and have a willingness to refocus and modify the aims as needed.

As an example, the following are aims of the first CMQCC Obstetric Hemorrhage Multi-Hospital Collaborative that began in October 2009 (Appendix E):

- **Aim 1:** Reduce the number of massive hemorrhages and the number of major complications from massive hemorrhage, including transfusions and hysterectomies, for all birthing women in participating hospitals by 75% by September 30, 2010.
- **Aim 2:** All collaborative participants develop and implement a multidisciplinary team response to every massive obstetric hemorrhage by September 30, 2010.

Complete a MAP-IT planning worksheet for your project (See Appendix F). The plan should focus on specific objectives that will help you achieve the overall aims of the project. Developing a plan for each objective facilitates communication and tracking of the project. The MAP-IT plan needs to be regularly reviewed and re-adjusted based on feedback obtained during the rapid cycles of change and small tests of improvement. In Appendix F is a sample of a completed MAP-IT Plan for the objective to quantify blood loss in order to recognize an obstetric hemorrhage.

Implement

STEP 4. IMPLEMENT STRATEGIES AND TACTICS TO MITIGATE BARRIERS

A. Implementation Strategies and Tactics

There are three broad types of implementation strategies the project team can utilize during implementation: 1) Education, 2) Data, and 3) Discourse or Communication

Within each strategy there are multiple implementation tactics that leaders can choose from. Tables 1, 2 and 3 below outline some specific strategies and tactics for improving obstetric hemorrhage practices.(9, 10)

Table 1. Examples of Implementation EDUCATION Tactics

Educational Tactics Examples	Definition and Examples Specific to OB Hemorrhage
Grand Rounds	<p><i>Physician educational sessions that are often held once a week.</i></p> <ul style="list-style-type: none"> • Hold a multi-disciplinary Obstetric Hemorrhage grand rounds using the standard toolkit slide set • Make a copy of the slides and put them in a binder in L&D for all staff to review
Classes or Conferences	<p><i>Formal educational sessions developed on specific topics.</i></p> <ul style="list-style-type: none"> • Develop a class and skills station on quantification of blood loss – use hospital specific products
Simulation Training	<p><i>Simulation training is education that allows clinicians to practice skills and knowledge through a fabricated situation that mimics a complicated situation that they will face and need to practice how to respond.</i></p> <ul style="list-style-type: none"> • Organize and run multi-disciplinary drills that help teams learn to follow the CMQCC Obstetric Hemorrhage checklist so they have a more organized and systematic approach to hemorrhage • This strategy also helps teams uncover hidden systems problems within their facility and get teams excited about finding solutions.
Competency Fairs, Tests, Learning Fairs, Return Demonstrations	<p><i>Clinicians demonstrate their knowledge of a new concept or demonstrate their ability to perform a clinical skill.</i></p> <ul style="list-style-type: none"> • Develop methods of measuring and tracking competency of team members for the quantification of blood loss • Be creative and make learning fun
On-Line Learning	<p><i>The use of the internet for the transfer of information.</i></p> <ul style="list-style-type: none"> • Provide clinicians with information on how to log-in to CMQCC’s website and have access to all of the CMQCC OB Hemorrhage resources: Table Chart, Check-list, Flow Chart, Compendium of Best Practices, and research articles

Table 2. Examples of Implementation DATA Tactics

Examples of Data Tactics	Definition and Examples Specific to OB Hemorrhage
Audit and feedback (group and individual)	<p><i>An examination of clinical records in order to gather specific pre-determined information. The information gathered are summarized and shared with the relevant group or individuals.</i></p> <ul style="list-style-type: none"> • Revise charting methods to track critical data • Develop a quality improvement data collection plan using audit tools (refer to Appendices E1-E5) • Develop a quality improvement measurement grid (refer to Appendix F)
Public Release of Data	<p><i>Details of care patterns and outcomes are reported openly to the community in such a way that anyone can access this information.</i></p> <ul style="list-style-type: none"> • Improve how obstetric hemorrhage is defined and coded to make it comparisons among hospitals more accurate

*Adapted from Bingham D. Measuring and increasing the effectiveness of the quality improvement implementation change practices of front-line maternity physician and nurse leaders, The University of North Carolina at Chapel Hill, 2009, 316 pages; AAT 3352932.

Table 3. Examples of Implementation COMMUNICATION Tactics

Examples of Communication Tactics	Definition
Meetings	<p><i>Group discussions, e.g., staff meetings.</i></p> <ul style="list-style-type: none"> • Regularly discuss the goals of the project and chart audit results at staff meetings and at other gatherings • Present project data to senior leadership at hospital wide meetings
One-to-One Discussions	<p><i>A discussion between a change leader and someone else that they are seeking to influence to change.</i></p> <ul style="list-style-type: none"> • Team members can promote the change by explaining the project goals during one-to-one discussions
Debriefs	<p><i>A team discussion that takes place as soon as possible after an emergency as possible.</i></p> <ul style="list-style-type: none"> • Complete a debrief form for every stage 2 and 3 obstetric hemorrhage emergency (refer to Appendix E.5)
Academic Detailing	<p><i>A review of relevant academic research by one leader meeting with one clinician at a time. A common tactic utilized by pharmaceutical representatives.</i></p>
Policy and Procedure	<p><i>A document utilized by a hospital to set and communicate clinical standards.</i></p> <ul style="list-style-type: none"> • Update policies and procedures
Reminders	<p><i>A method for helping someone remember to perform specific tasks, e.g., checklists, order set.</i></p> <ul style="list-style-type: none"> • Have blood loss markings added to the pouch on the under buttocks drape

Table 3. Examples of Implementation COMMUNICATION Tactics, *continued*

Examples of Communication Tactics	Definition
Newsletters	<i>A formal written update that is periodically distributed.</i>
Posters and Bulletin Boards	<i>A collection of data and information that are organized for display on poster board or a bulletin board.</i> <ul style="list-style-type: none"> • Develop posters and post on the unit.
Emails	<i>Electronic communications either one-to-one or by means of a “listserv” that is broadcast to many recipients.</i>
Rewards	<i>Something given in compensation for reaching a pre-determined goal, e.g., professional recognition, a bonus.</i> <ul style="list-style-type: none"> • Make learning fun – have some contests on assessing quantification of blood loss
Disciplinary Discussions	<i>A discussion that is held by someone in the position to give employee feedback and a formal review of performance in order to outline how current behavior do not meet required expectations of job performance.</i> <ul style="list-style-type: none"> • Follow-up with individuals who refuse to adopt the new practice standards

*Adapted from Bingham D. Measuring and increasing the effectiveness of the quality improvement implementation change practices of front-line maternity physician and nurse leaders, The University of North Carolina at Chapel Hill, 2009, 316 pages; AAT 3352932.

B. Potential Implementation Barriers

The most effective implementation plans include targeted strategies to mitigate potential and identified barriers. (10) Some types of barriers that have been identified are outlined in Table 4.

Table 4. Examples of Potential Implementation Barriers*

Leader Barriers	
Lack of Leader Knowledge to:	Design, plan, and implement Quality Improvement (QI)
	Perform QI data analysis
	Assess how to enhance their individual QI leadership abilities
Leader Attitudes (Beliefs and Assumptions) that Affect:	Topic selection
	QI topic goals
	Selection of QI implementation tactics
	Definitions of success
Leader Practices	Lack of clarity of QI project goals
	Backing down or stop trying
	Lack of time and other resources
	Inadequate practices to ensure sustainability, e.g., new hires, staff returning from vacation, leaves of absences
Clinician Barriers	
Lack of Clinician Knowledge:	About their own practices (lack adequate feedback)
	New or inexperienced
	Lack knowledge of the QI project
Clinician Attitudes (Beliefs and Assumptions):	Not persuaded to change
	Want autonomy
Clinician Practices:	Inertia – no motivation to change
	Forget
	Changes add more work or slow down usual work flow
Characteristics of the QI Project	
	Positive or negative effect(s) on clinician income or time
	Complexity of the QI project, e.g., how many groups' work flow is affected by the QI project changes
Implementation Climate/Culture	
	Type of hospital
	Amount of resources, e.g., workload, lack of organizational support for time or supplies needed to implement change
	Type of community or population of patients
	History of previous unsuccessful change attempts or minimal previous history of successful change attempts

*Adapted from Bingham D. Measuring and increasing the effectiveness of the quality improvement implementation change practices of front-line maternity physician and nurse leaders, The University of North Carolina at Chapel Hill, 2009, 316 pages; AAT 3352932.

Track

STEP 5. TRACK PROGRESS USING ESTABLISHED MEASURES

Essential components of measures include outcome, process and balance measures. When creating measures, teams should consider: What measures will be useful at your specific hospital based on your population of patients?

Below are some definitions of different types of measures.

- **Outcome Measures:** These measures tell you whether changes are actually leading to improvement – that is, helping to achieve the overall aim of reducing major complications of OB hemorrhage. Outcome measures answer questions like, “How many women had injuries?” and “How many women had markers that indicate potential morbidities (e.g., transfusions?)”. Outcome measures can sometimes be collected from administrative data.
- **Process Measures:** To affect the outcome measures of reducing major complications of OB hemorrhage, changes will be made to improve many core processes in the care system, as well as changes to improve the culture as it relates to safety. We will want to know if the parts/steps in the system are performing as planned. Measuring the results of these process changes will tell you if the changes are leading to an improved, safer system.
- **Balancing Measures:** Measures to make sure that changes to improve one part of the system are not causing new problems in other parts of the system. Balancing measures can also help teams to draw reasonable conclusions about the sustainability of the changes.

CMQCC developed Obstetric Hemorrhage quality improvement measurements for a multi-hospital collaborative. These measures are outlined in Appendix G. Hospital leaders can choose which of these measures are critical for tracking progress and whether they are achieving their aims and objectives.

The tracking phase can also utilize the analysis tools outlined under the assessment phase.

Repeat the MAP-IT Process

Evaluating results early and often at each stage of implementation—before, during and after—will guide decision-making for small tests of change. Small tests of change done over a short time frame and reviewed frequently provide necessary feedback to the team leaders. The feedback will help leaders re-adjust the MAP-IT Plan appropriately.

APPENDICES

APPENDIX A. SAMPLE HEMORRHAGE POLICY AND PROCEDURE

Obstetric Hemorrhage Care Guidelines: Sample Policy and Procedure

POLICY INDEX: O	Page 1 of X
POLICY TITLE: Obstetric Hemorrhage Care Guidelines	
DEPARTMENT AND USERS DISTRIBUTION: Maternal Child Health, Labor and Delivery, Emergency Department, Operating Room, Blood Bank, Intensive Care Unit, Post-Anesthesia Care Unit(s)	

Original Date of Issue: _____

Reviewed Date						
Revised Date						

PURPOSE

The purpose of this protocol is to provide guidelines for the optimal response of the multidisciplinary team in the event of obstetric hemorrhage. This protocol will also aid in recognizing patients at risk for hemorrhage and identifying stages of hemorrhage and primary treatment goals.

POLICY STATEMENTS

Optimal response to obstetric hemorrhage requires the coordination of effort of team members from multiple disciplines and departments.

- Obstetric unit, anesthesia department, blood bank, operating room, and other appropriate services work together to identify necessary system supports and processes for mounting an efficient and coordinated response to obstetric hemorrhage.
- Obstetric physicians, obstetric RNs, certified nurse midwives, anesthesiologists, and other appropriately qualified clinicians are authorized to mobilize the team to respond to an obstetric hemorrhage.
- The OB hemorrhage critical pack/cart are always kept stocked, not expired, and available for an emergency in all areas of the hospital where women are treated for OB hemorrhage. Note: the assignments for stocking and checking the cart need to be clearly delineated by each hospital. For example: medications will be kept together in an emergency packet in the pharmacy cart on the unit; the emergency medication packet will be maintained by pharmacy; the adult resuscitation cart or a separate resuscitation cart will be designed with an OB hemorrhage supply component.
- The Obstetric (OB) Hemorrhage general and massive policies and procedures will be updated at least every three years.

DEFINITIONS

General Hemorrhage: ≥ 500 ml blood loss for vaginal birth; ≥ 1000 ml blood loss for cesarean birth
Massive Hemorrhage: ≥ 1500 ml blood loss for any birth

MONITORING

Perform annual assessment of readiness to respond to an obstetric hemorrhage.

SUMMARY OF STAGES OF OBSTETRIC HEMORRHAGE AND PRIMARY TREATMENT GOALS

Prenatal Screening and Treatment:

Risk assessment
Aggressive treatment of anemia
Risk appropriate blood work on admission

Stage 0: Prevention and Recognition of OB Hemorrhage in All Births
Active Management of Third Stage Labor
Ongoing Quantitative Evaluation of Blood Loss
Ongoing Evaluation of Vital Signs

Stage 1: Cumulative Blood Loss >500 ml vaginal birth or >1000 ml cesarean birth **-OR-**
Vital Signs $>15\%$ change or HR ≥ 110 , BP $\leq 85/45$, O₂ sat $<95\%$ **-OR-**
Increased bleeding during recovery or postpartum
ACTIVATE HEMORRHAGE PROTOCOL, INITIATE PREPARATIONS, GIVE METHERGINE IM ONCE; IF NO RESPONSE, MOVE TO PROSTAGLANDINS (HEMABATE, CYTOTEC) (See Uterotonic Agent Information Table; Addendum A)

Stage 2: Continued Bleeding or Vital Sign instability and 1000-1500 ml cumulative blood loss
SEQUENTIALLY ADVANCE THROUGH MEDICATIONS AND PROCEDURES;
MOBILIZE HELP & BLOOD BANK SUPPORT;
KEEP AHEAD WITH VOLUME AND BLOOD PRODUCTS

Stage 3: Cumulative Blood Loss >1500 ml, >2 units PRBCs given, Vital Signs unstable or suspicion for Disseminated Intravascular Coagulopathy
ACTIVATE MASSIVE TRANSFUSION PROTOCOL AND INVASIVE SURGICAL APPROACHES TO CONTROL BLEEDING

PROCEDURES

Prenatal, Admission and Ongoing Risk Assessment

- Identify and prepare for patients with special considerations: Placenta Preview/Accreta, Bleeding Disorder, or those who decline blood products
- Screen and aggressively treat severe anemia: if oral iron fails, initiate IV Iron Sucrose Protocol ([Best Practice: Iron Sucrose Protocol](#)) to reach desired Hgb/Hct, especially for at-risk mothers

Admission Assessment & Planning

<p>Verify Type & Antibody Screen from prenatal record</p> <p>If not available,</p> <ul style="list-style-type: none"> <input type="checkbox"/> Order Type & Screen (lab will notify if 2nd clot needed for confirmation) <p>If prenatal or current antibody screen positive (if not low level anti-D from Rho-GAM),</p> <ul style="list-style-type: none"> <input type="checkbox"/> Type & Crossmatch 2 units PRBCs <p>All other patients,</p> <ul style="list-style-type: none"> <input type="checkbox"/> Send Clot to blood bank 	<p>Evaluate for Risk Factors (see below)</p> <p>If medium risk:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Order Type & Screen <input type="checkbox"/> Review Hemorrhage Protocol <p>If high risk:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Order Type & Crossmatch 2 units PRBCs <input type="checkbox"/> Review Hemorrhage Protocol <input type="checkbox"/> Notify OB Anesthesia <p>Identify women who may decline transfusion</p> <ul style="list-style-type: none"> <input type="checkbox"/> Notify OB provider for plan of care <input type="checkbox"/> Early consult with OB anesthesia <input type="checkbox"/> Review Consent Form
--	---

Ongoing Risk Assessment

<ul style="list-style-type: none"> <input type="checkbox"/> Evaluate for development of additional risk factors in labor: <ul style="list-style-type: none"> • Prolonged 2nd Stage labor • Prolonged oxytocin use • Active bleeding • Chorioamnionitis • Magnesium sulfate treatment <input type="checkbox"/> Increase Risk level (see below) and convert to Type & Screen or Type & Crossmatch <input type="checkbox"/> Treat multiple risk factors as High Risk
--

Admission Hemorrhage Risk Factor Evaluation

Low (Clot only)	Medium (Type and Screen)	High (Type and Cross)
No previous uterine incision	Prior cesarean birth(s) or uterine surgery	Placenta previa, low lying placenta,
Singleton pregnancy	Multiple gestation	Suspected placenta accreta or percreta
≤4 previous vaginal births	>4 previous vaginal births	Hematocrit <30 AND other risk factors
No known bleeding disorder	Chorioamnionitis	Platelets <100,000
No history of PPH	History of previous PPH	Active bleeding (greater than show) on admit
	Large uterine fibroids	Known coagulopathy
	Estimated fetal weight greater than 4 kg	
	Morbid obesity (BMI >35)	

*If admitted patients are started on magnesium sulfate they are at higher risk of postpartum hemorrhage.

PROCEDURES, CONTINUED

STAGE 0

Prevention & Recognition of Hemorrhage during all births

Active Management of Third Stage of Labor

1. Administer Oxytocin infusion: 10-20 Units/1000 ml solution for women with IV access. Note that the dosage and rates should be clearly specified by each hospital.
 - a. Titrate infusion rate to uterine tone.
 - b. Use 10 units IM for women without IV access.
 - c. **Do not give oxytocin as IV push**
2. Provide vigorous fundal massage for at least 15 seconds

Ongoing Quantitative Measurement of Blood Loss at all Births

1. Assess blood loss at birth, prior to delivery of the placenta whenever possible.
2. Reassess cumulative blood loss after delivery of the placenta
3. Use formal methods to assess blood loss:
 - a. Use graduated under-buttock drapes
 - b. Weigh blood soaked materials on gram scale (1 gm = 1ml)
 - i. Subtract known dry weight of materials
 - ii. Use a hemorrhage report or [Early Warning Chart \(National Health Survey, NHS\)](#)

**NOTE: if a dry chux is used to protect scale from blood-soaked material, ZERO the scale after placing dry chux and prior to placing saturated item(s).*

Ongoing Evaluation of Vital Signs and Clinical Triggers

STAGE 1

Cumulative Blood Loss >500 ml vaginal birth or >1000 ml C/S -OR- Vital Signs >15% change or HR ≥110, BP ≤85/45, O2 sat <95% -OR- Increased bleeding during recovery or postpartum

Interventions:

Follow [Obstetric hemorrhage care guidelines checklist](#) to mobilize response, act to mitigate bleeding, and move sequentially through treatment.

Evaluate patient response to interventions:

1. If the patient is stable following Stage 1 interventions then perform increased postpartum surveillance.

STAGE 2

Proceed to STAGE 2 *for any of the following* when cumulative blood loss is <1500 mL:

1. Continued bleeding
2. Continued vital sign instability

Evaluate patient response to interventions:

1. If stabilized during Stage 2 (<1500 ml cumulative blood loss) then perform increased postpartum surveillance
-

STAGE 3

Proceed to STAGE 3 if cumulative blood loss >1500 mL OR:

1. >2 units PRBCs administered
2. Unstable vital signs after stage 2 interventions
3. Suspicion of DIC

Evaluate patient response to interventions:

1. If stabilized during Stage 3 (cumulative blood loss >1500 ml) then perform increased postpartum surveillance, consult with intensivist and/or transfer to ICU
-

Do not delay other interventions while waiting for response to medication(s).

Do not wait for laboratory values to initiate transfusions:

1. Transfuse based on clinical signs and patient response.
2. Transfuse aggressively with a high ratio of Fresh Frozen Plasma to PRBCs for massive hemorrhage (>1500 mL cumulative blood loss); key is high ratio of FFP to PRBC
 - Either 6:4:1 PRBCs:FFP:Platelets
 - Or 4:4:1 PRBCs:FFP:Platelets

COMMUNICATION and DOCUMENTATION

1. Verbally acknowledge actions you will take and orders received.
2. Provide ongoing updates about patient's status with other departments.
3. Record intake and output records.

REFERENCES and RELATED DOCUMENTS:

CMQCC Obstetric Hemorrhage Tool-Kit, April 2009, www.cmqcc.org.

1. CMQCC OB Hemorrhage Task Force: Care Guidelines and Compendium of Best Practices, [OB Hemorrhage Care Guidelines Checklist](#): use the checklist to help think through possible etiologies and anticipate next steps and to identify Risk Factors: Prenatal, Admission and Ongoing Assessment
2. Lyndon, A., et al, [Ongoing Quantitative Measurement of Blood Loss](#) at all births
3. Casper, L., Lee, R., [Carts, Kits and Trays](#)
4. Gregory, K., et al, [Definitions, Early Recognition, and Rapid Response Using Triggers](#)

Addendum A

UTEROTONIC AGENTS for POSTPARTUM HEMORRHAGE						
Drug	Dose	Route	Frequency	Side Effects	Contraindications	Storage
Pitocin® (Oxytocin) 10 units/ml	10-40 units per 1000 ml, rate titrated to uterine tone	IV infusion	Continuous	Usually none Nausea, vomiting, hyponatremia ("water intoxication") with prolonged IV admin. ↓ BP and ↑ HR with high doses, esp IV push	Hypersensitivity to drug	Room temp
Methergine® (Methylergonivine) 0.2mg/ml	0.2 mg	IM (not given IV)	-Q 2-4 hours -If no response after first dose, it is unlikely that additional doses will be of benefit	Nausea, vomiting Severe hypertension, esp. with rapid administration or in patients with HTN or PIH	Hypertension, PIH, Heart disease Hypersensitivity to drug Caution if multiple doses of ephedrine have been used, may exaggerate hypertensive response w/possible cerebral hemorrhage	Refrigerate Protect from light
Hemabate® (15-methyl PG F2a) 250mcg/ml	250 mcg	IM or intra-myometrial (not given IV)	-Q 15-90 min -Not to exceed 8 doses/24 hrs -If no response after several doses, it is unlikely that additional doses will be of benefit.	Nausea, vomiting, Diarrhea Fever (transient), Headache Chills, shivering Hypertension Bronchospasm	Caution in women with hepatic disease, asthma, hypertension, active cardiac or pulmonary disease Hypersensitivity to drug	Refrigerate
Cytotec® (Misoprostol) 100 or 200mcg tablets	800-1000mcg	Per rectum (PR)	One time	Nausea, vomiting, diarrhea Shivering, Fever (transient) Headache	Rare Known allergy to prostaglandin Hypersensitivity to drug	Room temp

APPENDIX B: CMQCC OB HEMORRHAGE CARE GUIDELINES CHECKLIST

CMQCC CALIFORNIA MATERNAL QUALITY CARE COLLABORATIVE **Obstetric Hemorrhage Care Guidelines: Checklist Format** version 1.4

Prenatal Assessment & Planning

- Identify and prepare for patients with special considerations:** Placenta Previa/Accreta, Bleeding Disorder, or those who Decline Blood Products
- Screen and aggressively treat severe anemia:** if oral iron fails, initiate IV Iron Sucrose Protocol to reach desired Hgb/Hct, especially for at risk mothers.

Admission Assessment & Planning

Ongoing Risk Assessment

Verify Type & Antibody Screen from prenatal record
If not available,
 Order Type & Screen (lab will notify if 2nd clot needed for confirmation)
If prenatal or current antibody screen positive (if not low level anti-D from Rho-GAM),
 Type & Crossmatch 2 units PRBCs
All other patients,
 Send Clot to blood bank

Evaluate for **Risk Factors** (see below)
If medium risk:
 Order Type & Screen
 Review Hemorrhage Protocol
If high risk:
 Order Type & Crossmatch 2 units PRBCs
 Review Hemorrhage Protocol
 Notify OB Anesthesia
Identify women who may decline transfusion
 Notify OB provider for plan of care
 Early consult with OB anesthesia
 Review Consent Form

- Evaluate for development of additional risk factors in labor:**
 - Prolonged 2nd Stage labor
 - Prolonged oxytocin use
 - Active bleeding
 - Chorioamnionitis
 - Magnesium sulfate treatment
- Increase Risk level** (see below) **and convert to Type & Screen or Type & Crossmatch**
- Treat multiple risk factors as High Risk**

Admission Hemorrhage Risk Factor Evaluation		
Low (Clot only)	Medium (Type and Screen)	High (Type and Crossmatch)
No previous uterine incision	Prior cesarean birth(s) or uterine surgery	Placenta previa, low lying placenta
Singleton pregnancy	Multiple gestation	Suspected Placenta accreta or percreta
≤4 previous vaginal births	>4 previous vaginal births	Hematocrit <30 AND other risk factors
No known bleeding disorder	Chorioamnionitis	Platelets <100,000
No history of PPH	History of previous PPH	Active bleeding (greater than show) on admit
	Large uterine fibroids	Known coagulopathy
	Estimated fetal weight greater than 4 kg	
	Morbid obesity (BMI >35)	

STAGE 0: All Births: Prevention & Recognition of OB Hemorrhage

- Active Management of Third Stage**
- Oxytocin infusion: 10-20 units oxytocin/1000ml solution titrate infusion rate to uterine tone; or 10 units IM; do not give oxytocin as IV push
 - Vigorous **fundal** massage for at least 15 seconds
- Ongoing Quantitative Evaluation of Blood Loss**
- Using formal methods, such as graduated containers, visual comparisons and weight of blood soaked materials (**1gm = 1ml**)
- Ongoing Evaluation of Vital Signs**

If: Cumulative Blood Loss >500ml vaginal birth or >1000ml C/S -OR- Vital signs >15% change or HR ≥110, BP ≤85/45, O2 sat <95% -OR- Increased bleeding during recovery or postpartum, proceed to STAGE 1

APPENDIX B: CHECKLIST, continued

STAGE 1: OB Hemorrhage Cumulative Blood Loss >500ml vaginal birth or >1000ml C/S -OR- Vital signs >15% change or HR ≥110, BP ≤85/45, O2 sat <95% -OR- Increased bleeding during recovery or postpartum		
MOBILIZE	ACT	THINK
Primary nurse, Physician or Midwife to: <input type="checkbox"/> Activate OB Hemorrhage Protocol and Checklist Primary nurse to: <input type="checkbox"/> Notify obstetrician (in-house and attending) <input type="checkbox"/> Notify charge nurse <input type="checkbox"/> Notify anesthesiologist	Primary nurse: <input type="checkbox"/> Establish IV access if not present, at least 18 gauge Increase IV fluids rates (Lactated Ringers preferred) and increase Oxytocin rate (500 mL/hour of 10-40 units/1000mL solution); Titrate Oxytocin infusion rate to uterine tone <input type="checkbox"/> Continue vigorous fundal massage <input type="checkbox"/> Administer Methergine 0.2 mg IM per protocol (if not hypertensive); give once, if no response, move to alternate agent; if good response, may give additional doses q 2 hr <input type="checkbox"/> Vital Signs, including O2 sat & level of consciousness (LOC) q 5 minutes <input type="checkbox"/> Weigh materials, calculate and record cumulative blood loss q 5-15 minutes <input type="checkbox"/> Administer oxygen to maintain O2 sats at >95% <input type="checkbox"/> Empty bladder: straight cath or place Foley with urimeter <input type="checkbox"/> Type and Crossmatch for 2 units Red Blood Cells STAT (if not already done) <input type="checkbox"/> Keep patient warm Physician or midwife: <input type="checkbox"/> Rule out retained Products of Conception, laceration, hematoma Surgeon (if cesarean birth and still open) <input type="checkbox"/> Inspect for uncontrolled bleeding at all levels, esp. broad ligament, posterior uterus, and retained placenta	Consider potential etiology: <ul style="list-style-type: none"> • Uterine atony • Trauma/Laceration • Retained placenta • Amniotic Fluid Embolism • Uterine Inversion • Coagulopathy • Placenta Accreta • Uterine Rupture Once stabilized: Modified Postpartum management with increased surveillance
If: Continued bleeding or Continued Vital Sign instability, and <1500 mL cumulative blood loss proceed to STAGE 2		

UTEROTONIC AGENTS for POSTPARTUM HEMORRHAGE						
Drug	Dose	Route	Frequency	Side Effects	Contraindications	Storage
Pitocin® (Oxytocin) 10 units/ml	10-40 units per 1000 ml, rate titrated to uterine tone	IV infusion	Continuous	Usually none Nausea, vomiting, hyponatremia ("water intoxication") with prolonged IV admin. ↓ BP and ↑ HR with high doses, esp IV push	Hypersensitivity to drug	Room temp
Methergine® (Methylergonivine) 0.2mg/ml	0.2 mg	IM (not given IV)	-Q 2-4 hours -If no response after first dose, it is unlikely that additional doses will be of benefit	Nausea, vomiting Severe hypertension, esp. with rapid administration or in patients with HTN or PIH	Hypertension, PIH, Heart disease Hypersensitivity to drug Caution if multiple doses of ephedrine have been used, may exaggerate hypertensive response w/possible cerebral hemorrhage	Refrigerate Protect from light
Hemabate® (15-methyl PG F2a) 250mcg/ml	250 mcg	IM or intra-myometrial (not given IV)	-Q 15-90 min -Not to exceed 8 doses/24 hrs -If no response after 3 doses, it is unlikely that additional doses will be of benefit.	Nausea, vomiting, Diarrhea Fever (transient), Headache Chills, shivering Hypertension Bronchospasm	Caution in women with hepatic disease, asthma, hypertension, active cardiac or pulmonary disease Hypersensitivity to drug	Refrigerate
Cytotec® (Misoprostol) 100 or 200mcg tablets	800-1000mcg	Per rectum (PR)	One time	Nausea, vomiting, diarrhea Shivering, Fever (transient) Headache	Rare Known allergy to prostaglandin Hypersensitivity to drug	Room temp

California Maternal Quality Care Collaborative (CMQCC): Hemorrhage Taskforce (2009) visit: www.CMQCC.org for details

STAGE 2: OB Hemorrhage Continued bleeding or Vital Sign instability, and <1500 mL cumulative blood loss		
MOBILIZE	ACT	THINK
<p>Primary nurse (or charge nurse):</p> <ul style="list-style-type: none"> <input type="checkbox"/> Call obstetrician to bedside <input type="checkbox"/> Call Anesthesiologist <input type="checkbox"/> Activate Response Team: <p>PHONE #: _____</p> <ul style="list-style-type: none"> <input type="checkbox"/> Notify Blood bank of hemorrhage; order products as directed <p>Charge nurse:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Notify Perinatologist or 2nd OB <input type="checkbox"/> Initiate OB Hemorrhage Record <input type="checkbox"/> If selective embolization, call in Interventional Radiology Team and second anesthesiologist <input type="checkbox"/> Notify nursing supervisor <input type="checkbox"/> Assign single person to communicate with blood bank <input type="checkbox"/> Call medical social worker or assign other family support person 	<p>Team leader (OB physician):</p> <ul style="list-style-type: none"> <input type="checkbox"/> Additional uterotonic medication: Hemabate 250 mcg IM [if not contraindicated] OR Misoprostol 800-1000 mcg PR <ul style="list-style-type: none"> o Can repeat Hemabate up to 3 times every 20 min; (note-75% respond to first dose) <p>Do not delay other interventions (see right column) while waiting for response to medications</p> <ul style="list-style-type: none"> <input type="checkbox"/> Bimanual uterine massage <input type="checkbox"/> Move to OR (if on postpartum unit, move to L&D or OR) <input type="checkbox"/> Order 2 units PRBCs and bring to the bedside <input type="checkbox"/> Order labs STAT (CBC/Plts, Chem 12, PT/aPTT, Fibrinogen, ABG) <input type="checkbox"/> Transfuse PRBCs based on clinical signs and response, do not wait for lab results <p>Primary nurse:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Establish 2nd large bore IV, at least 18 gauge. Maintain adequate fluid volume with Lactated Ringers and adequate uterine tone with oxytocin infusion <input type="checkbox"/> Assess and announce Vital Signs and cumulative blood loss q 5-10 minutes <input type="checkbox"/> Set up blood administration set and blood warmer for transfusion <input type="checkbox"/> Administer meds, blood products and draw labs, as ordered <input type="checkbox"/> Keep patient warm <p>Second nurse (or charge nurse):</p> <ul style="list-style-type: none"> <input type="checkbox"/> Place Foley with urimeter (if not already done) <input type="checkbox"/> Obtain portable light and OB procedure tray or Hemorrhage cart <input type="checkbox"/> Obtain blood products from the Blood Bank <input type="checkbox"/> Assist with move to OR (if indicated) <p>Blood Bank:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Determine availability of thawed plasma, fresh frozen plasma, and platelets; initiate delivery of platelets if not present on-site <input type="checkbox"/> Consider thawing 2 FFP (takes 30 min), use if transfusing >2 units PRBCs <input type="checkbox"/> Prepare for possibility of massive hemorrhage 	<p>Sequentially advance through procedures and other interventions based on etiology:</p> <p>Vaginal birth</p> <p>If trauma (vaginal, cervical or uterine):</p> <ul style="list-style-type: none"> • Visualize and repair <p>If retained placenta:</p> <ul style="list-style-type: none"> • D&C <p>If uterine atony or lower uterine segment bleeding:</p> <ul style="list-style-type: none"> • Intrauterine Balloon <p>If above measures unproductive:</p> <ul style="list-style-type: none"> • Selective embolization (Interventional Radiology if available & adequate experience) <p>C-section:</p> <ul style="list-style-type: none"> • Uterine hemostatic suture, e.g., B-Lynch Suture, O'Leary, Multiple Squares • Intrauterine Balloon <p>If Uterine Inversion:</p> <ul style="list-style-type: none"> • Anesthesia and uterine relaxation drugs for manual reduction <p>If Amniotic Fluid Embolism:</p> <ul style="list-style-type: none"> • Maximally aggressive respiratory, vasopressor and blood product support <p>If vital signs are worse than estimated or measured blood loss: possible uterine rupture or broad ligament tear with internal bleeding; move to laparotomy</p> <p>Once stabilized: Modified Postpartum management with increased surveillance</p>
<p>Re-Evaluate Bleeding and Vital Signs If cumulative blood loss >1500ml, >2 units PRBCs given, VS unstable or suspicion for DIC, proceed to STAGE 3</p>		


STAGE 3: OB Hemorrhage Cumulative blood loss >1500ml, >2 units PRBCs given, VS unstable or suspicion for DIC		
MOBILIZE	ACT	THINK
<p>Nurse or Physician:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Activate Massive Hemorrhage Protocol <p>PHONE #: _____</p> <p>Charge Nurse or designee:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Notify advanced Gyn surgeon (e.g. Gyn Oncologist) <input type="checkbox"/> Notify adult intensivist <input type="checkbox"/> Call-in second anesthesiologist <input type="checkbox"/> Call-in OR staff <input type="checkbox"/> Reassign staff as needed <input type="checkbox"/> Call-in supervisor, CNS, or manager <input type="checkbox"/> Continue OB Hemorrhage Record (In OR, anesthesiologist will assess and document VS) <input type="checkbox"/> If transfer considered, notify ICU <p>Blood Bank:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Prepare to issue additional blood products as needed – stay ahead 	<p>Establish team leadership and assign roles</p> <p>Team leader (OB physician + OB anesthesiologist, anesthesiologist and/or perinatologist and/or intensivist):</p> <ul style="list-style-type: none"> <input type="checkbox"/> Order Massive Hemorrhage Pack (RBCs + FFP + 1 pheresis pack PLTS—see note in right column) <input type="checkbox"/> Move to OR if not already there <input type="checkbox"/> Repeat CBC/PLTS, Chem 12, PT/aPTT, Fibrinogen, ABG STAT q 30-60 min <p>Anesthesiologist (as indicated):</p> <ul style="list-style-type: none"> <input type="checkbox"/> Arterial blood gases <input type="checkbox"/> Central hemodynamic monitoring <input type="checkbox"/> CVP or PA line <input type="checkbox"/> Arterial line <input type="checkbox"/> Vasopressor support <input type="checkbox"/> Intubation <p>Primary nurse:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Announce VS and cumulative measured blood loss q 5-10 minutes <input type="checkbox"/> Apply upper body warming blanket if feasible <input type="checkbox"/> Use fluid warmer and/or rapid infuser for fluid & blood product administration <input type="checkbox"/> Apply sequential compression stockings to lower extremities <input type="checkbox"/> Circulate in OR <p>Second nurse and/or anesthesiologist:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Continue to administer meds, blood products and draw labs, as ordered <p>Third Nurse (or charge nurse):</p> <ul style="list-style-type: none"> <input type="checkbox"/> Recorder 	<ul style="list-style-type: none"> • Selective Embolization (IR) • Interventions based on etiology not yet completed • Prevent hypothermia, Acidemia <p>Conservative or Definitive Surgery:</p> <ul style="list-style-type: none"> • Uterine Artery Ligation • Hysterectomy <div style="border: 1px solid black; padding: 5px; margin: 10px 0;"> <p style="text-align: center;">For Resuscitation: Aggressively Transfuse Based on Vital Signs, Blood Loss</p> <p style="text-align: center;">KEY: HIGH RATIO of FFP to RBC Either: 6:4:1 PRBCs: FFP: Platelets Or: 4:4:1 PRBCs: FFP: Platelets</p> </div> <p>Unresponsive Coagulopathy:</p> <ul style="list-style-type: none"> • After 8-10 units PRBCs and coagulation factor replacement may consider risk/benefit of rFactor VIIa <p>Once Stabilized: Modified Postpartum Management; consider ICU</p>

BLOOD PRODUCTS	
<p>Packed Red Blood Cells (PRBC) <i>(approx. 35-40 min. for crossmatch—assuming no sample is in the lab and assuming no antibodies are present)</i> Transfuse O Negative blood if you cannot wait</p>	<p>Best first-line product for blood loss 1 unit = 450ml volume If antibody positive, may take 1-24 hrs. for crossmatch 1 unit=450 ml volume and typically increases Hct by 3%</p>
<p>Fresh Frozen Plasma (FFP) <i>(approx. 35-45 min. to thaw for release)</i></p>	<p>Highly desired if >2 units PRBCs given, or for prolonged PT, aPTT >1.5x control 1 unit = 180ml volume and typically increases Fibrinogen by 10mg/dL</p>
<p>Platelets (PLTS) <i>Local variation in time to release (may need to come from regional blood bank)</i></p>	<p>Priority for women with Platelets <50,000 Single-donor Apheresis unit (= 6 units of platelet concentrates) provides 40-50k transient increase in platelets</p>
<p>Cryoprecipitate (CRYO) <i>(approx. 35-45 min. to thaw for release)</i></p>	<p>Priority for women with Fibrinogen levels <80 10 unit pack typically raises Fibrinogen 80-100mg/dL Best for DIC with low fibrinogen and don't need volume replacement Caution: 10 units come from 10 different donors, so infection risk is proportionate.</p>

APPENDIX C. CMQCC OBHEMORRHAGE CARE GUIDELINES FLOW CHART



OBSTETRIC HEMORRHAGE CARE SUMMARY: FLOW CHART FORMAT



California Maternal Quality Care Collaborative (CMQCC), Hemorrhage Taskforce (2009) visit: www.CMQCC.org for details
This project was supported by Title V funds received from the State of California Department of Public Health, Center for Family Health; Maternal, Child and Adolescent Health Division

APPENDIX D. CMQCC OB HEMORRHAGE CARE GUIDELINES TABLE CHART



Obstetric Hemorrhage Care Summary: Table Chart Format version 1.4

	Assessments	Meds/Procedures	Blood Bank
Stage 0	Every woman in labor/giving birth		
<i>Stage 0 focuses on risk assessment and active management of the third stage.</i>	<ul style="list-style-type: none"> Assess every woman for risk factors for hemorrhage Ongoing quantitative evaluation of blood loss on every birth 	Active Management 3rd Stage: <ul style="list-style-type: none"> Oxytocin IV infusion or 10u IM Fundal Massage-vigorous, <u>15 seconds min.</u> 	<ul style="list-style-type: none"> If Medium Risk: T&Scr If High Risk: T&C 2 U If Positive Antibody Screen (prenatal or current, exclude low level anti-D from RhoGam): T&C 2 U
Stage 1	Blood loss: >500 ml vaginal <u>or</u> >1000 ml Cesarean, <u>or</u> VS changes (by >15% <u>or</u> HR ≥110, BP ≤85/45, O2 sat <95%)		
<i>Stage 1 is short: activate hemorrhage protocol, initiate preparations and give Methergine IM.</i>	<ul style="list-style-type: none"> Activate OB Hemorrhage Protocol and Checklist Notify Charge nurse, Anesthesia Provider VS, O2 Sat q5' Calculate cumulative blood loss q5-15' Weigh bloody materials Careful inspection <u>with good exposure</u> of vaginal walls, cervix, uterine cavity, placenta 	<ul style="list-style-type: none"> IV Access: at least 18gauge Increase IV fluid (LR) and Oxytocin rate, and repeat fundal massage Methergine 0.2mg IM (if not hypertensive) May repeat if good response to first dose, BUT otherwise move on to 2nd level uterotonic drug (see below) Empty bladder: straight cath or place foley with urimeter 	<ul style="list-style-type: none"> T&C 2 Units PRBCs (if not already done)
Stage 2	Continued bleeding with total blood loss under 1500ml		
<i>Stage 2 is focused on sequentially <u>advancing</u> through medications and procedures, mobilizing help and Blood Bank support, and keeping ahead with volume and blood products.</i>	OB back to bedside (if not already there) <ul style="list-style-type: none"> Extra help: 2nd OB, Rapid Response Team (per hospital), assign roles VS & cumulative blood loss q 5-10 min Weigh bloody materials Complete evaluation of vaginal wall, cervix, placenta, uterine cavity Send additional labs, including DIC panel If in Postpartum: Move to L&D/OR Evaluate for special cases: <ul style="list-style-type: none"> -Uterine Inversion -Amn. Fluid Embolism 	2nd Level Uterotonic Drugs: <ul style="list-style-type: none"> Hemabate 250 mcg IM <u>or</u> Misoprostol 800-1000 mcg PR 2nd IV Access (at least 18gauge) Bimanual massage Vaginal Birth: (typical order) <ul style="list-style-type: none"> Move to OR Repair any tears D&C: r/o retained placenta Place intrauterine balloon Selective Embolization (Interventional Radiology) Cesarean Birth: (still intra-op) (typical order) <ul style="list-style-type: none"> Inspect broad lig, posterior uterus and retained placenta B-Lynch Suture Place intrauterine balloon 	<ul style="list-style-type: none"> Notify Blood Bank of OB Hemorrhage Bring 2 Units PRBCs to bedside, transfuse per clinical signs – do not wait for lab values Use blood warmer for transfusion Consider thawing 2 FFP (takes 35+min), use if transfusing >2u PRBCs Determine availability of additional RBCs and other Coag products
Stage 3	Total blood loss over 1500ml, <u>or</u> >2 units PRBCs given <u>or</u> VS unstable <u>or</u> suspicion of DIC		
<i>Stage 3 is focused on the Massive Transfusion protocol and invasive surgical approaches for control of bleeding.</i>	<ul style="list-style-type: none"> Mobilize team <ul style="list-style-type: none"> -Advanced GYN surgeon -2nd Anesthesia Provider -OR staff -Adult Intensivist Repeat labs including coags and ABG's Central line Social Worker/ family support 	<ul style="list-style-type: none"> Activate Massive Hemorrhage Protocol Laparotomy: <ul style="list-style-type: none"> -B-Lynch Suture -Uterine Artery Ligation -Hysterectomy Patient support <ul style="list-style-type: none"> -Fluid warmer -Upper body warming device -Sequential compression stockings 	Transfuse Aggressively Massive Hemorrhage Pack <ul style="list-style-type: none"> Near 1:1 PRBC:FFP 1 PLT pheresis pack per 6units PRBCs Unresponsive Coagulopathy: After 10 units PRBCs <u>and</u> full coagulation factor replacement: may consider rFactor VIIa

This Project was supported by Title V funds received from the State of California, Department of Public Health, Center for Family Health; Maternal, Child and Adolescent Health Division

APPENDIX E. CMQCC OBSTETRIC HEMORRHAGE QUALITY IMPROVEMENT COLLABORATIVE

CMQCC initiated a statewide Obstetric Hemorrhage Quality Improvement (QI) Collaborative in October 2009. The OB Hemorrhage QI Collaborative is using the Institute for Healthcare Improvement (IHI) Breakthrough Series Model for Improvement, which includes intensive expert and peer mentoring. The Expert Panel, with input from the collaborating hospitals, developed specific goals or aims for the 12-month initiative. The aims guided the development of measures that assess improvement over the life of the collaborative and beyond.

Aims and measures from this Collaborative are presented as a guide, but can be changed to meet facilities' needs.

Refer to the appendix for checklists, flowcharts, audit tools and additional documents to use during implementation and change.

APPENDIX E.1. CMQCC IN HOSPITAL AUDIT TOOL: RISK ASSESSMENT FOR OB HEMORRHAGE

Topic: Risk Assessment for obstetric hemorrhage is documented in the chart at admission.

Goal: 100% of women are assessed for risk of obstetric hemorrhage on admission by [date].

Instructions: Audit 20 *randomly selected* charts per month (10 Vaginal, 10 cesarean).

Risk Assessment is documented in the chart	MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery					
	Yes <input type="checkbox"/> No <input type="checkbox"/>					
Risk Assessment is documented in the chart	MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery					
	Yes <input type="checkbox"/> No <input type="checkbox"/>					
Risk Assessment is documented in the chart	MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery					
	Yes <input type="checkbox"/> No <input type="checkbox"/>					
Risk Assessment is documented in the chart	MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery					
	Yes <input type="checkbox"/> No <input type="checkbox"/>					

Total number of audited charts with:

Numerator: _____ Risk Assessment Documented in Chart

Denominator: _____ Total Number of Charts Audited

APPENDIX E.2. CMQCC IN HOSPITAL AUDIT TOOL: ACTIVE MANAGEMENT OF THIRD STAGE LABOR

Topic: Active Management of Third Stage Labor* is documented in the chart at admission.

Goal: 100% of women are assessed for risk of obstetric hemorrhage on admission by [date].

Instructions: Audit 20 *randomly selected* charts per month (10 Vaginal, 10 cesarean). Select “All are recorded” check box ***only if all 3 elements*** of Active Management* were documented in the chart.

*Active Management of Third Stage Labor is defined as including ALL of the following:

1. Oxytocin (IV or IM) at delivery of shoulders or delivery of placenta (identify when administered)
2. Fundal Massage for 15 seconds minimum
3. Gentle Cord Traction

MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery <input type="checkbox"/> ALL are recorded (check if Yes): • Oxytocin IV/IM <i>Administered at delivery of (check one):</i> <input type="checkbox"/> Shoulders -OR- <input type="checkbox"/> Placenta • Fundal Massage (≥ 15 sec) • Cord Traction	MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery <input type="checkbox"/> ALL are recorded (check if Yes): • Oxytocin IV/IM <i>Administered at delivery of (check one):</i> <input type="checkbox"/> Shoulders -OR- <input type="checkbox"/> Placenta • Fundal Massage (≥ 15 sec) • Cord Traction	MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery <input type="checkbox"/> ALL are recorded (check if Yes): • Oxytocin IV/IM <i>Administered at delivery of (check one):</i> <input type="checkbox"/> Shoulders -OR- <input type="checkbox"/> Placenta • Fundal Massage (≥ 15 sec) • Cord Traction	MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery <input type="checkbox"/> ALL are recorded (check if Yes): • Oxytocin IV/IM <i>Administered at delivery of (check one):</i> <input type="checkbox"/> Shoulders -OR- <input type="checkbox"/> Placenta • Fundal Massage (≥ 15 sec) • Cord Traction	MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery <input type="checkbox"/> ALL are recorded (check if Yes): • Oxytocin IV/IM <i>Administered at delivery of (check one):</i> <input type="checkbox"/> Shoulders -OR- <input type="checkbox"/> Placenta • Fundal Massage (≥ 15 sec) • Cord Traction
MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery <input type="checkbox"/> ALL are recorded (check if Yes): • Oxytocin IV/IM <i>Administered at delivery of (check one):</i> <input type="checkbox"/> Shoulders -OR- <input type="checkbox"/> Placenta • Fundal Massage (≥ 15 sec) • Cord Traction	MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery <input type="checkbox"/> ALL are recorded (check if Yes): • Oxytocin IV/IM <i>Administered at delivery of (check one):</i> <input type="checkbox"/> Shoulders -OR- <input type="checkbox"/> Placenta • Fundal Massage (≥ 15 sec) • Cord Traction	MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery <input type="checkbox"/> ALL are recorded (check if Yes): • Oxytocin IV/IM <i>Administered at delivery of (check one):</i> <input type="checkbox"/> Shoulders -OR- <input type="checkbox"/> Placenta • Fundal Massage (≥ 15 sec) • Cord Traction	MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery <input type="checkbox"/> ALL are recorded (check if Yes): • Oxytocin IV/IM <i>Administered at delivery of (check one):</i> <input type="checkbox"/> Shoulders -OR- <input type="checkbox"/> Placenta • Fundal Massage (≥ 15 sec) • Cord Traction	MR# <input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery <input type="checkbox"/> ALL are recorded (check if Yes): • Oxytocin IV/IM <i>Administered at delivery of (check one):</i> <input type="checkbox"/> Shoulders -OR- <input type="checkbox"/> Placenta • Fundal Massage (≥ 15 sec) • Cord Traction

Total number of audited charts with:

Numerator: _____ **All 3 elements of** Active Management of Third Stage Labor Documented in Chart

Denominator: _____ Total Number of Charts Audited

APPENDIX E.3. METHODS FOR DEVELOPING TRAINING AND TOOLS FOR QUANTITATIVE MEASUREMENT OF BLOOD LOSS

Recommended methods for ongoing quantitative measurement of blood loss:

1. Formally estimate blood loss by recording percent (%) saturation of blood soaked items with the use of visual cues such as pictures/posters to determine blood volume equivalence of saturated/blood soaked pads, chux, etc.
2. Formally measure blood loss by weighing blood soaked pads/chux
3. Formally measure blood loss by collecting blood in graduated measurement containers

Quantifying blood loss by weighing (see images at right and below)


- Establish dry weights of common items
- Standardize use of pads
- Build weighing of pads into routine practice
- Develop worksheet for calculations

Quantifying blood loss by measuring (see image below right)


- Use graduated collection containers (C/S and vaginal deliveries)
- Account for other fluids (amniotic fluid, urine, irrigation)

Training Tools

Posters



18 X 18 inch Dry Lap Sponges



- 25 ml saturates about 50% area
- 50 ml saturates about 75% area
- 75 ml saturates entire surface
- 100 ml will saturate and drip


Establish Dry Weights

Dry Weights		Procedure
Item	Weight in Grams	
Standard Bundle (2 lg chux, 1 ice pack peripad, 2 small peripads)	398	<ul style="list-style-type: none"> • Weigh all bloody items in grams • Subtract dry weights in grams • Remaining weight in grams = ml blood loss <p>1 gram = 1 ml</p>
Small Chux (16 in X 24 in)	22	
Large Chux (24 in X 34 in)	98	
Large Peripad (peach backing)	26	
Small Peripad (from OB Pack)	15	
Ice Pack Peripad	172	
Cloth Towel (blue)	88 - 115	
Vag Packing (from OB Pack)	18	
Ray-tec Sponge	4	

**Posters
Pocket Cards**



Used with kind permission of Bev VanderWal, CNS



APPENDIX E.4. CMQCC IN HOSPITAL AUDIT TOOL: CUMULATIVE BLOOD LOSS AND QUANTITATIVE MEASUREMENT METHODS



QI Chart Review Data Collected by: _____

Date Collected: _____

IN HOSPITAL AUDIT TOOL: CUMULATIVE BLOOD LOSS AND QUANTITATIVE MEASUREMENT METHODS

Topic: Cumulative blood loss is recorded in patient chart (in mls) during labor and delivery, until status is routine postpartum and patient is physiologically stable.

Goal: 100% of birthing mothers have on-going cumulative quantification of blood loss by September 30, 2010.

Instructions: Audit 20 *randomly selected* charts per month (10 Vaginal, 10 Cesarean; identify method of delivery below). Identify whether blood loss was evaluated; identify if blood loss was evaluated by visual estimation only; identify if blood loss was quantitatively evaluated using one or more of three recommended methods.*

Submit Cumulative Blood Loss Audit Data to CMQCC monthly using EXTRANET (do not submit medical record numbers)

Four methods for Ongoing cumulative blood loss measurement:

1. Estimating blood loss by visual estimation methods only

***Recommended methods:**

2. Formally estimate blood loss by recording percent (%) saturation of blood soaked items with the use of visual cues such as pictures/posters to determine blood volume equivalence

3. Formally measure blood loss by weighing blood soaked pads/chux

4. Formally measure blood loss by collecting blood in graduated measurement containers

MR #	<input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery
Cumulative blood loss is recorded in patient chart in milliliters during labor and delivery and until status changes to routine postpartum and patient is physiologically stable.	<input type="checkbox"/> Measurement NOT recorded in chart <input type="checkbox"/> Estimated with Visual Cues <i>Only</i> If you select Visual Estimation Only, do not select methods below
*Select the Measurement Method(s) Recorded in Patient Chart	Select All that Apply: <input type="checkbox"/> Formally estimated by % saturation <input type="checkbox"/> Formally measured by weighing <input type="checkbox"/> Formally measured by collection

MR #	<input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery
Cumulative blood loss is recorded in patient chart in milliliters during labor and delivery and until status changes to routine postpartum and patient is physiologically stable.	<input type="checkbox"/> Measurement NOT recorded in chart <input type="checkbox"/> Estimated with Visual Cues <i>Only</i> If you select Visual Estimation Only, do not select methods below
*Select the Measurement Method(s) Recorded in Patient Chart	Select All that Apply: <input type="checkbox"/> Formally estimated by % saturation <input type="checkbox"/> Formally measured by weighing <input type="checkbox"/> Formally measured by collection

MR #	<input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery
Cumulative blood loss is recorded in patient chart in milliliters during labor and delivery and until status changes to routine postpartum and patient is physiologically stable.	<input type="checkbox"/> Measurement NOT recorded in chart <input type="checkbox"/> Estimated with Visual Cues <i>Only</i> If you select Visual Estimation Only, do not select methods below
*Select the Measurement Method(s) Recorded in Patient Chart	Select All that Apply: <input type="checkbox"/> Formally estimated by % saturation <input type="checkbox"/> Formally measured by weighing <input type="checkbox"/> Formally measured by collection

MR #	<input type="checkbox"/> Vaginal Delivery <input type="checkbox"/> Cesarean Delivery
Cumulative blood loss is recorded in patient chart in milliliters during labor and delivery and until status changes to routine postpartum and patient is physiologically stable.	<input type="checkbox"/> Measurement NOT recorded in chart <input type="checkbox"/> Estimated with Visual Cues <i>Only</i> If you select Visual Estimation Only, do not select methods below
*Select the Measurement Method(s) Recorded in Patient Chart	Select All that Apply: <input type="checkbox"/> Formally estimated by % saturation <input type="checkbox"/> Formally measured by weighing <input type="checkbox"/> Formally measured by collection

Total number of audited charts with Blood Loss:

Numerator 1: _____ Not Measured

Numerator 2: _____ Estimated with Visual Cues Only

Numerator 3: _____ Measured using one or more of three recommended formal measurements (estimate by % saturation; measure by weight; measure by collection)

Denominator (for all 3 numerators above) _____ Total Number of Charts Audited

APPENDIX E.5. CMQCC OBSTETRIC HEMORRHAGE TEAM DE-BRIEFING FORM

Topic: The de-brief form provides an opportunity for maternity service teams to review then document sequence of events, successes and barriers to a swift and coordinated response to obstetric hemorrhage.

Goal: De-brief completed in 100% of all obstetric hemorrhages that progress to Stage 2 or 3. All de-briefs have at least Primary RN, and Primary MD who participates in the de-briefing session.

Instructions: Complete as soon as possible, but no later than 24 hours after any Stage 2 or 3 hemorrhages. During de-brief, obtain input from participants (all or as many as possible).

Stage 2 or 3 hemorrhages are defined as bleeding that continues after administration of IV or IM Oxytocin, vigorous fundal massage, emptied bladder and Methergine 0.2 mg IM.

PRINT THIS FORM DOUBLE-SIDED; PATIENT STAMP ON BACK OF FORM WILL NOT BE FAXED WITH COMPLETED FORM. - Attach additional pages with notes as needed Were the following medications, procedures or blood products used? (Check if yes, check all that apply)

Medications

- High dose misoprostol (800-1000 mcg)
- Carboprost tromethamine (Hemobate)

Blood Volume/Options

- Invasive hemodynamic monitoring
- Blood warmer
- Rapid fluid infuser (level one machine)
- Blood cell salvage machine (cell saver)
- Factor VIIa (non-standard treatment)

Procedures

- Intrauterine balloons
- B-Lynch suture
- Uterine artery ligation
- Uterine artery embolization
- Non-pneumatic Anti-shock Garments (NASG; non-standard treatment)

COMMENTS about medications, procedures, or blood products:

- Primary MD participated in de-brief? (Check if yes)
- Primary RN participated in de-brief? (Check if yes)

Post-hemorrhage, the patient required...

- (Check if yes, check all that apply)
- Intubation
 - Pressors
 - Admission to ICU
 - Central Line
 - Arterial Line
 - Admission to higher acuity unit (e.g., PACU)

Volume of blood lost: _____ mls

Method of Blood Loss Measurement (Check all that apply)

- Visually Estimated Only
- Formal Estimate using Posters/Pictures
- Formal Measure by weight
- Formal Measure by volume collection

Blood Product Transfusion Ratios - Active Hemorrhage Treatment and Resuscitation Period (~the first 4-6 hours PP)

Units of PRBCs: _____ Units of FFP: _____

Units of Platelets: _____ Units of Cryo: _____

Thinking about how the obstetric hemorrhage was managed...

Identify what went well (Check if yes, describe)

- Communication went well
- Teamwork went well
- Leadership went well
- Decision-making went well
- Assessing the situation went well
- Other

Briefly describe:

Identify opportunities for improvement: “non-human factors” (Check if yes, describe)

- Delay in blood products availability
- Equipment issues
- Medications issues

- Inadequate support (in-unit or other areas of the hospital)
- Delays in transporting the patient (within the hospital or to another facility)
- Other

Briefly describe:

Identify opportunities for improvement: “human factors” (Check if yes, describe)

- Communication needed improvement
- Teamwork needed improvement
- Leadership needed improvement
- Decision-making needed improvement
- Assessing needed improvement
- Other

Briefly describe:

APPENDIX F. CMQCC MAP-IT PLANNING WORKSHEET, SAMPLE WORKSHEET

Change Project MAP-IT Worksheet – Blank Form

MAP-IT Action Plan for: _____(Hospital Name)
Date Created: _____ Developed by: _____

Aims Statement or Objective: *By (month)____ (day)____ (year)____ we will quantify blood loss on all women who give birth at our hospital.*

M: Mobilize

A: Assess

P: Plan

I: Implement

T: Track

First Cycle Due Date: _____

Guidry, M., Vischi, T., Han, R., & Passons, O. *Healthy people in healthy communities: A community planning guide using healthy people 2010*. Washington, D.C.: U.S. Department of Health and Human Services. The Office of Disease Prevention and Health Promotion.
<http://www.healthypeople.gov/Publications/HealthyCommunities2001/default.htm>.

APPENDIX F. CMQCC MAP-IT PLANNING WORKSHEET, SAMPLE WORKSHEET

**SAMPLE of COMPLETED
OB Hemorrhage Change Project MAP-IT Planning Worksheet**

MAP-IT Action Plan for: California's Best Medical Center _____ (Hospital Name)
Date Created: _____ Developed by: OB Hemorrhage QI Team

Aims Statement or Objective: *By (month)____ (day)____ (year)____ we will quantify blood loss on all women who give birth at our hospital.*

M: Mobilize

Get your change team together on a regular basis. Have the perspectives of a Hospital Administrator, Labor & Delivery leader and staff, Anesthesia, Blood Bank, Emergency Department, etc. represented on your team so that they can develop a plan for each of their departments.

A: Assess

Complete the Fishbone cause and effect diagram on quantification of blood loss. Complete a process flow chart on the current state of blood loss calculation and accuracy.

P: Plan

1. Review the policy and procedures to determine how they need to be revised.
2. Raise awareness among the staff of how inaccurate the blood loss measurements currently are. A. In a room on or near the unit set up a "Quantification of Blood Loss Skills Lab". The skills lab can have multiple stations where there are pads, chux, etc. stained and saturated with red fluid. Have the clinicians guess the amounts of fluid on each of the items. Post the results for everyone to see. Make it fun! B. Have team members make rounds on all shifts and ask the clinicians who were at a delivery to quantify the blood loss. Weigh the items saturated with blood and compare the actual amounts to the estimated amounts.
3. Revise the chart forms to include method of determination of blood loss.
4. Identify who will be responsible to accomplish each of the above items.

I: Implement

1. Revise the policy and procedure to outline how blood loss will be quantified. Have a select group of staff test the draft procedure to see how the procedure needs to be refined and adjusted and what barriers are identified. Eliminate the barriers identified and revise the procedure as needed.
2. Assess all staff on their ability to accurately assess blood loss. Randomly compare estimated blood loss to actual blood loss.
3. Revise charting forms. Trial the new forms. Re-adjust as needed.
4. Keep each other apprised of the work that is accomplished and any barriers identified. Provide each other support for the implementation efforts.

T: Track

After implementing the above steps, assess the number of staff who quantify blood loss. Develop a new plan and implement that plan. Track progress and re-adjust the plan again and again until the goal is met and the change has become "how things are done around here."

First Cycle Due Date: _____

Guidry, M., Vischi, T., Han, R., & Passons, O. *Healthy people in healthy communities: A community planning guide using healthy people 2010*. Washington, D.C.: U.S. Department of Health and Human Services. The Office of Disease Prevention and Health Promotion. <http://www.healthypeople.gov/Publications/HealthyCommunities2001/default.htm>

APPENDIX G. CMQCC QUALITY IMPROVEMENT COLLABORATIVE OBSTETRIC HEMORRHAGE MEASUREMENT GRID

**CMQCC Quality Improvement Collaborative
Obstetric Hemorrhage Measurement Grid**

Purpose of the Measurement Grid:

The measurement grid outlines the measures to be collected over the 12-month life of the OB Hemorrhage multi-hospital collaborative. The grid includes the specific parameters for each measure.

Measurement for Improvement:

CMQCC multi-hospital improvement collaboratives are about making hospital systems safer for patients. Measurements play an important role. Always remember that measurements should be designed to accelerate improvement, not slow it down.

Timeline for Measurement:

The measurement grid is broken down into three categories: Outcome, Process and Balancing Measures. Teams may also develop additional measures based on the issues that are of most interest and importance to their hospital and patient population needs. Only the measures in the grid below will be submitted to CMQCC.

Outcome Measures: These measures tell you whether changes are actually leading to improvement – that is, helping to achieve the overall aim of reducing major complications of OB hemorrhage. Outcome measures answer questions like, “How many women had injuries?” and “How many women had markers that indicate potential morbidities (e.g., transfusions?)”. Outcome measures can sometimes be collected from administrative data.

Process Measures: To affect the outcome measures of reducing major complications of OB hemorrhage, changes will be made to improve many core processes in the care system, as well as changes to improve the culture as it relates to safety. We will want to know if the parts/steps in the system are performing as planned. Measuring the results of these process changes will tell you if the changes are leading to an improved, safer system.

Balancing Measures: We will use these measures to make sure that changes to improve one part of the system are not causing new problems in other parts of the system. Balancing measures can also help us to draw reasonable conclusions about the sustainability of the change

OB Hemorrhage Multi-Hospital Change Collaborative

Aim 1: Reduce the number of massive hemorrhages and the number of major complications from massive hemorrhage, including transfusions and hysterectomies, for all birthing women in participating hospitals by 75% by September 30, 2010.

Aim 2: All collaborative participants develop and implement a multidisciplinary team response to every massive obstetric hemorrhage by September 30, 2010.

OB HEMORRHAGE TOOLKIT

Measure	Calculation	Data Collection Plan/Sample Size	Data Collection Method	Goal
<p>Aim 1. Blood Product Transfusions Measures A, B. REQUIRED Measure Types: Outcome</p>				
<p>REQUIRED A.1. Percent of women (who gave birth ≥ 20 0/7 weeks gestation) who were transfused with any blood product during the birth admission</p> <p>A.2. Total units of each type of blood product (PRBCs, Platelets, Plasma/FFP, Cryo) transfused during birth admissions per total births</p> <p>REQUIRED B. Percent of women (who gave birth ≥ 20 0/7 weeks gestation) who were transfused with ≥ 5 units PRBCs during the birth admission NOTE: B is a subset of A.1</p>	<p>A.1. Numerator: Number of women (who gave birth ≥ 20 0/7 weeks gestation) who were transfused with any blood product during the birth admission per month.</p> <p>Denominator: Total Number of Births (≥ 20 0/7 weeks gestation) per month</p> <p>A.2. Numerators: Total units of each type of blood product (PRBCs, Platelets, Plasma/FFP, Cryo) transfused during birth admissions per month. Select each blood product within the series from the drop-down menu:</p> <ul style="list-style-type: none"> • Series 1*: PRBCs units/month • Series 2: Platelets units/month • Series 3: Plasma/FFP units/month • Series 4: Cryo units/month <p><i>*Series are categories of data within a single measure</i></p> <p>Denominator: Total Number of Births (≥ 20 0/7 weeks gestation) per month</p> <p>B) Numerator: Number of women (who gave birth ≥ 20 0/7 weeks gestation) who were transfused with ≥ 5 units PRBCs during the birth admission per month.</p> <p>Denominator: Total Number of Births (≥ 20 0/7 weeks gestation) per month.</p>	<p>Hospital Baseline: Collect A1, A2, B for January 9, 2009 to September 30, 2009</p> <p>Concurrent: Report monthly data as defined (starting with October 2009)</p> <p>OPTIONAL Additional Measurement: Number of Births/Days between occurrences of massive hemorrhage</p>	<p>Blood Transfusion: data from internal source such as blood bank data, patient charts, medical records, Electronic Medical Record (EMR), etc.</p> <p>If available: Blood loss data recorded in patient record or delivery log.</p> <p>ICD-9 Procedure Code for transfusions: 99.0</p> <p>CPT Code: 36430: Transfusion, blood or blood components</p> <p><i>Note that these codes typically do not accurately identify transfusions. We recommend obtaining data from the Blood Bank when possible.</i></p> <p>Data entered to Extranet</p> <p>De-brief Form: For women who experience Stage 2 or 3 hemorrhage, identify units of PRBCs, Platelets, Plasma/FFP, Cryo (for each woman) on the De-brief Form</p>	<p>Reduce the number of major complications from massive hemorrhage, including transfusions, for all birthing women in participating hospitals by 75% by September 30, 2010.</p>

Measure	Calculation	Data Collection Plan/Sample Size	Data Collection Method	Goal
Aim 1. Peripartum Hysterectomies Measure C. REQUIRED Measure Type: Outcome				
<p>C. Rate of peripartum hysterectomies in women (who gave birth ≥ 20 0/7 weeks gestation) per 1000 births (hysterectomy performed during birth admission) stratified by risk of Placenta Previa and/or Placenta Accreta/percreta</p>	<p>C. Numerator: Number of peripartum hysterectomies (performed during birth admission) in women (who gave birth ≥ 20 0/7 weeks gestation) per month stratified by:</p> <ul style="list-style-type: none"> • Series 1*: Women with Placenta Previa and/or Placenta Accreta/Percreta • Series 2: Women without Placenta Previa and/or Placenta Accreta/Percreta <p><i>*Series are categories of data within a single measure</i></p> <p>Denominator: Total Number of Births (≥ 20 0/7 weeks gestation) per month</p> <p>Risk Stratification/Adjustment: Women who had a hysterectomy and placenta previa and/or accreta/percreta are reported separately from women who had a hysterectomy (and NO placenta previa/accreta/percreta)</p> <p>Annotate** for each hysterectomy:</p> <ol style="list-style-type: none"> a) Indication for hysterectomy b) Number of prior cesarean sections c) Number of Days Post-Delivery (Days = 0 if procedure done on day of delivery) 	<p>Hospital Baseline: Number of peripartum hysterectomies (performed during birth admission) in women who gave birth ≥ 20 0/7 weeks gestation Between 1/1/09 to 9/30/09</p> <p>Concurrent: Report monthly data as defined</p>	<p>Peripartum Hysterectomy: Data Collection from internal source such as EMR, medical records, or other method determined by each site</p> <p>ICD-9 Procedure Codes 68.3 Subtotal abdominal hysterectomy 68.39 Other and unspecified subtotal abdominal hysterectomy 68.4 Total abdominal hysterectomy 68.49 Other and unspecified total abdominal hysterectomy</p> <p>CPT Codes 59525 Cesarean Hysterectomy 58150 Hysterectomy Total/Partial (Use Post-Partum or with Vaginal) 59160 D&C after delivery</p> <p>Data entered to Extranet **Annotation is available in the Data Entry fields for this measure; identify a), b), and c) (from Calculation column) for each patient in the Annotation Field</p>	<p>Reduce the number of major complications from massive hemorrhage, including peripartum hysterectomies, for all birthing women in participating hospitals by 75% by September 30, 2010.</p>

Measure	Calculation	Data Collection Plan/Sample Size	Data Collection Method	Goal
<p>Aim 1. Quantification of Blood Loss Measures D, E, F. REQUIRED Deliverables 1, 2, 3. REQUIRED Measure Types: Outcome and Process</p>				
<p>D. Percent of Audited Charts in which quantification and documentation of blood loss is performed (during and after all births until immediate recovery status changes to routine postpartum care and woman is physiologically stable) using one or more of three preferred methods:</p> <ol style="list-style-type: none"> Formally estimate blood loss by recording percent (%) saturation of blood soaked items with the use of visual cues such as pictures/posters to determine blood volume equivalence of saturated/blood soaked pads, chux, etc. Formally measure blood loss by weighing blood soaked pads/chux Formally measure blood loss by collecting blood in graduated measurement containers <p>Process Measures:</p> <p>E. % Non-MD clinicians and staff who are educated about Cumulative Blood Loss and Quantitative Measurement Methods</p>	<p>D. Numerator: Number of charts per month where on-going quantification is:</p> <ul style="list-style-type: none"> Series 1: <u>NOT</u> recorded Series 2: Recorded <u>using visualization only</u> Series 3: Recorded using one of three preferred formal methods (1, 2, or 3 in Measure Column) <p>Denominator: Number of charts audited per month</p> <p>E. Numerator: Number of non-MD clinicians, (e.g., RNs, midwives) and staff (e.g., clerks, aides) who are educated about Cumulative Blood Loss and Quantitative Measurement Methods Audit Tool per month</p> <p>Denominator: Number of Non-MD clinicians and staff who care for</p>	<p>Audit 20 randomly selected charts per month (10 vaginal births and 10 cesarean births)</p>	<p>Chart Review: Refer to sample audit tool entitled: "In Hospital Audit Tool: Cumulative Blood Loss and Quantitative Measurement Methods"</p> <p>Data entered to Extranet</p>	<p>100% of birthing women will have on-going cumulative quantification of blood loss by September 30, 2010.</p>

<p>F. % MDs (e.g., obstetricians, anesthesiologists) who are educated about Cumulative Blood Loss and Quantitative Measurement Methods</p> <p>Deliverables:</p> <ol style="list-style-type: none"> 1. Sign-off report among Labor & Delivery and Post-partum MD and Nursing staff includes whether a woman had a Stage 2 or Stage 3 hemorrhage 2. Blood loss is measured until the woman's immediate recovery status changes to routine postpartum care and woman is physiologically stable 3. Documentation forms are updated for #1, #2 	<p>women giving birth in your facility per month</p> <p>F. Numerator: Number of MDs (e.g., obstetricians, anesthesiologists) who are educated about Cumulative Blood Loss and Quantitative Measurement Methods per month</p> <p>Denominator: Number of MDs who care for women giving birth in your facility per month</p> <ol style="list-style-type: none"> 1. Deliverable: Submit in Extranet when completed 2. Deliverable: Submit in Extranet when completed 3. Deliverable: Submit in Extranet when completed 	<ol style="list-style-type: none"> 1. Deliverable: Due 5/1/2010 2. Deliverable: Due 5/1/2010 3. Deliverable: Due 5/1/2010 		
--	---	---	--	--

Measure	Calculation	Data Collection Plan/Sample Size	Data Collection Method	Goal
Aim 1. OB Hemorrhage Risk Assessment on Admission Measure G. RECOMMENDED Measure Type: Process				
Percent of women (from audited charts) who are assessed for risk of obstetric hemorrhage on admission	<p>Numerator: Among the audited charts, the number of women admitted to Labor and Delivery whose risk of obstetric hemorrhage assessment is recorded in the medical record</p> <p>Denominator: Total number of admission charts audited per month</p>	<p>Suggestion: Audit 20 randomly selected charts per month (10 vaginal births and 10 cesarean births)</p>	Chart Review Determined by individual hospital. Refer to Risk Assessment Audit Tool. Data entered to Extranet	100% of women are assessed for risk of obstetric hemorrhage on admission by September 30, 2010
Aim 1. Active Management of Third Stage Measure H. RECOMMENDED Measure Type: Process				
Percent of women (from audited charts) who receive Active Management of Third Stage Labor	<p>Numerator: Total number of women who receive Active Management of Third Stage Labor including</p> <ul style="list-style-type: none"> • Oxytocin (IV or IM) at delivery of shoulders or delivery of placenta (identify when administered) • Fundal Massage for 15 seconds minimum • Gentle Cord traction <p><i>Note: Need all three to be considered Active Management</i></p> <p>Denominator: Total number of charts audited per month</p>	<p>Suggestion: Audit 20 randomly selected charts of women who gave birth vaginally per month</p>	Chart review Determined by individual hospital. Refer to Active Management of Third Stage Labor Audit Tool. Data entered to Extranet	100% of women giving birth will receive active management of labor by September 30, 2010

Measure	Calculation	Data Collection Plan/Sample Size	Data Collection Method	Goal
Aim 2. Policies and Procedures and Drills Measure I, J, K, L. REQUIRED Deliverables 4, 5, 6, 7, 8. REQUIRED Measure Type: Process				
<p>Deliverables:</p> <p>4. General and massive hemorrhage policies and procedures (P&P) were updated after April 30, 2009.</p> <p>5. Identify roles and multi-disciplinary team responders for stage 1, 2, and 3 hemorrhages</p> <p>6. Determine and implement the most desirable method for maintaining accessibility to the needed OB hemorrhage supplies</p> <p>Measures</p> <p>I. % Non-MD clinicians and staff who are educated to the hemorrhage P&Ps</p>	<p>4. Deliverable: Submit in Extranet when completed; Date completed; date received by CMQCC</p> <p>5. Deliverable: Roles defined for stage 1, 2, or 3 hemorrhages: Submit in Extranet when completed; date completed; date received by CMQCC</p> <p>6. Deliverable: Provide the emergency supply maintenance plan to CMQCC: Submit in Extranet when completed; date completed; date received by CMQCC</p> <p>I. Numerator: Number of non-MD clinicians, (e.g., RNs, midwives) and staff (e.g., clerks, aides) who receive education on the hemorrhage P&Ps per month</p> <p>Denominator: Number of Non-MD clinicians and staff who are in the pool of possible responders per month (defined by hospital)</p>	<p>4. Determined by hospital, Due: 2/1/2010</p> <p>5. Determined by hospital, Due: 3/1/2010</p> <p>6. Determined by hospital, Due: 2/1/2010</p> <p>I. Determined by hospital</p>	<p>Hospital records and forms</p>	<p>100% of the collaborative participating hospitals will meet 100% of the P&Ps and drills measurements by September 30, 2010</p>

<p>J. % MDs (e.g., obstetricians, anesthesiologists) who are educated to the hemorrhage P&Ps</p> <p>Deliverables</p> <p>7. Create drills tailored to your hospital P&Ps and responder roles</p> <p>8. After Deliverables 4-7 are completed, run 1 multi-disciplinary drill per month for four consecutive months (two on night/evening shift and two on day shift) to identify system and process improvement opportunities. After each drill complete a drill de-brief form</p> <p>Measure</p> <p>K. % Non-MD clinicians and staff who are involved drill de-brief discussions</p>	<p>J. Numerator: Number of MDs (e.g., obstetricians, anesthesiologists) who receive education on the hemorrhage P&Ps per month</p> <p>Denominator: Number of MDs who are in the pool of possible responders per month (defined by hospital)</p> <p>7. Deliverable: Drill scenarios created: Submit in Extranet when completed; date completed; date received by CMQCC</p> <p>8. Deliverable: Drills are performed: Submit in Extranet when completed; dates completed. Maintain a list of problems identified by the drills and document how and when the problem is resolved. Submit the list to CMQCC.</p> <p>K. Numerator: Number of non-MD clinicians (e.g., RNs, midwives) and staff (e.g., clerks, aides) who are involved in drill de-brief discussions per month</p>	<p>J. Determined by hospital</p> <p>7. Determined by hospital, Due: 5/1/2010</p> <p>8. Determined by hospital Due: 8/1/2010</p> <p>K. Determined by hospital</p>		
---	---	--	--	--

<p>L. % MDs (e.g., obstetricians, anesthesiologists) who are involved in drill de-brief discussions</p>	<p>Denominator: All Non-MD clinicians and staff who are in the pool of possible de-brief participants per month (defined by hospital)</p> <p>L. Numerator: Number of MDs (e.g., obstetricians, anesthesiologists) who are involved in drill de-brief discussions per month</p> <p>Denominator: Number of MDs who are in the pool of possible de-brief participants per month (defined by hospital)</p>	<p>L. Determined by hospital</p>		
---	---	----------------------------------	--	--

Measure	Calculation	Data Collection Plan/Sample Size	Data Collection Method	Goal
<p>Aim 2. De-Briefing after OB Hemorrhages</p>				
<p>Measures M, N, O. REQUIRED</p>				
<p>Measure Type: Process</p>				
<p>M. Frequency of de-brief sessions that occurred after any hemorrhage with ≥ 500 mls for vaginal births and ≥ 1000 mls for cesarean section births that advanced beyond stage 1 to stage 2 or 3 hemorrhages.</p> <p>Note: Stage 2 or 3 are defined as hemorrhages with bleeding that continues after the patient received IV or IM Oxytocin, vigorous fundal massage, emptied bladder,</p>	<p>M. Numerator: Number of de-brief forms submitted to CMQCC</p> <p>Denominator: Number of hemorrhages each month that required interventions, treatments, procedures outlined in Stage 2 or 3 of the CMQCC OB Hemorrhage checklist</p> <p>Recommendation: Completion of de-brief is encouraged to occur</p>	<p>M. RN who took care of the patient leads the de-briefing session and fills out de-briefing form</p>	<p>M. The CMQCC “Obstetric Hemorrhage Team De-Briefing Form” or any form that captures the elements contained on this form</p> <p>Fax de-brief form to CMQCC</p> <p>Data entered to Extranet (for Measures M, N, O)</p>	<p>De-brief completed in 100% of all obstetric hemorrhages that progress to Stage 2 or 3</p> <p>All de-briefs have at least one MD who participates in the de-briefing session.</p> <p>Improve communication, teamwork, use of equipment and overall management by de-briefing after every stage 2 and 3 hemorrhages.</p> <p>Identify barriers to and</p>

<p>and Methergine 0.2mg IM</p> <p>N. Percent of de-briefs during which at least one primary MD participated (as checked on de-brief form)</p> <p>O. Percent of de-briefs during which at least one primary RN participated (as checked on de-brief form)</p>	<p>immediately after the patient is stabilized, but no later than 24 hours after event.</p> <p>Numerator: Number of Primary MDs who participated in de-brief</p> <p>Denominator: Total number of de-brief forms completed</p> <p>O. Numerator: Number of Primary RNs who participated in de-brief</p> <p>Denominator: Total number of de-brief forms completed</p>	<p>Determined by hospital</p> <p>Determined by hospital</p>		<p>solutions for:</p> <ul style="list-style-type: none"> a. Communication; Teamwork; Leadership; Decision-making; Assessment (situational) b. Delays in blood product availability; Equipment issues; Medications issues; In-unit (and other) Support; Delays in patient transport
--	--	---	--	--

REFERENCES

1. Clark S, Miller D, Belford M, Dildy G, Frye D, Meyers J. Neonatal and maternal outcomes associated with elective term delivery. *American Journal of Obstetrics and Gynecology*. 2009;200:156.e1-.e4.
2. Lu MC, Fridman M, Korst LM, et al. Variations in the incidence of postpartum hemorrhage across hospitals in California. *Maternal Child Health Journal*. 2005 September;9(3):297-306.
3. Kuklina E, Meikle, S., Jamieson, D., Whiteman, M., Barfield, W., Hillis, S., Posner, S. Severe Obstetric Morbidity in the US, 1998-2005. *Obstetrics and Gynecology*. 2009;113:293-9.
4. Guidry M, Vischi T, Han R, Passons O. Healthy people in healthy communities: A community planning guide using healthy people 2010. US Department of Health and Human Services, The Office of Disease Prevention and Health Promotion <http://www.healthypeople.gov/Publications/HealthyCommunities2001/default.htm>. 2010.
5. Eckes G. Six Sigma for Everyone Hoboken, NJ: John Wiley & Sons. 2003.
6. Brussee W. Statistics for six sigma made easy! New York: McGraw-Hill; 2004.
7. Carey RG, Lloyd RC. Measuring quality improvement in healthcare: A guide to statistical process control applications. Milwaukee, Wisconsin: American Society for Quality; 2001.
8. Haugehy D. Pareto Analysis Step by Step <http://www.ginque.com/index.php/projmgmt/505-pareto-analysis-step-by-step>. Accessed March 31, 2010.
9. Bingham D. Measuring and increasing the effectiveness of the quality improvement implementation change practices of front-line maternity physician and nurse leaders. The University of North Carolina at Chapel Hill, 2009, 316 pages; AAT 3352932; 2009. p. 316.
10. Bingham D, Main E. Effective implementation strategies and tactics for leading change on maternity units. *Journal of Neonatal and Perinatal Nursing*. 2010 Accepted for publication;24(1):32-42.