



# STAT

Place X in box if STAT Medication Order.  
(If STAT box is marked, a new order sheet is required for future orders).

**Generic substitution and therapeutic interchange are authorized unless prohibited by the prescriber according to hospital policy**

**PATIENT STATUS:**  Admit to Antepartum Bed  Birthing Suites Bed

**TIME OF INDUCTION:**  Morning  Evening

**DIAGNOSIS:** \_\_\_\_\_

**NURSING CARE AND TREATMENTS:**

Last vaginal exam and confirmed vertex on \_\_\_\_\_ by \_\_\_\_\_ MD/CNM/PA  
(within 48 hours of induction start time)

Bishop Score  0-4  5-9  >10

Parameter Score	0	1	2	3
Position	Posterior	Intermediate	Anterior	-
Consistency	Firm	Intermediate	Soft	-
Effacement	0-30%	40-50%	60-70%	80%
Dilation	<1	1-2 cm	2-3 cm	>3cm
Fetal Position	-3	-2	-1, 0	+1, +2

Consent for vaginal delivery or cesarean section by Dr. \_\_\_\_\_

Notify MD of Admission

Initiate Labor and Delivery Admission Orders **(COMPLETE AND TRANSMIT TO PHARMACY)**

Epidural \_\_\_\_\_

GBBS protocol if GBBS positive: Penicillin 5 million units IV initially, then 3 million units IV every 4 hours until delivery

Antibiotic for GBBS Protocol for Penicillin Allergy \_\_\_\_\_ IV every \_\_\_\_\_ hours until delivery

Initiate Antepartum Cervical Ripening Orders **(COMPLETE AND TRANSMIT TO PHARMACY)**

\_\_\_\_\_

**If Being Induced for Preeclampsia**

Clean Catch Routine Urine

Cath Routine Urine

PEP Panel

**If Gestational Diabetic**

Diet Control Gestational Diabetic FSBS \_\_\_\_\_

Insulin Dependent Gestational Diabetic start Intrapartum Insulin Protocol **(COMPLETE AND TRANSMIT TO PHARMACY)**

Insulin Dependent Diabetic \_\_\_\_\_

**IV'S AND MEDICATIONS:**

**Type of Induction:**

**Cervical Ripening**

Misoprostol (Cytotec) 25 mcg intra-vaginally every 4 hours inserted by MD or RN

Misoprostol (Cytotec) 50 mcg intra-vaginally every 4 hours inserted by MD

Dinoprostone (Cervidil) 10 mg intra-vaginally inserted by MD

Foley bulb 30 mL to be placed by MD

**Pitocin**

Pitocin 30 units /500 mL IV titrate per Regular Protocol

Pitocin 30 units /500 mL IV titrate per Progressive Protocol

Start Pitocin per above 4 hours after last Cytotec or immediately after Cervidil removal

Begin Pitocin per above \_\_\_\_\_ hours after initial antibiotics for GBBS protocol

\_\_\_\_\_

**SEE PHYSICIAN ORDER SHEET FOR ADDITIONAL ORDERS**

**DATE:** \_\_\_\_\_ **TIME:** \_\_\_\_\_ **SIGNATURE:** \_\_\_\_\_ **M.D.**



FMOBG0010B

**OB INDUCTION ORDERS**